



HAL
open science

Exponential ergodicity of a degenerate age-size piecewise deterministic process

Ignacio Madrid

► **To cite this version:**

Ignacio Madrid. Exponential ergodicity of a degenerate age-size piecewise deterministic process. 2022.
hal-03885141v2

HAL Id: hal-03885141

<https://polytechnique.hal.science/hal-03885141v2>

Preprint submitted on 7 Dec 2022 (v2), last revised 18 Mar 2024 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exponential ergodicity of a degenerate age-size piecewise deterministic process

Ignacio Madrid*

Abstract

We study the long-time behaviour of the first-moment semigroup of a non conservative piecewise deterministic measure-valued stochastic process with support on \mathbb{R}_+^2 driven by a deterministic flow between random jump times, with a transition kernel which has a degenerate form. Using a Doob h -transform where the function h is taken as an eigenfunction of the associated generator, we can bring ourselves back to the study of a conservative process whose exponential ergodicity is proven via Harris' Theorem. Particular attention is given to the proof of Doeblin minoration condition. The main difficulty is the degeneracy of one of the two variables, and the deterministic dependency between the two variables, which make it no trivial to uniformly bound the expected value of the trajectories with respect to a non-degenerate measure in a two-dimensional space, which is particularly hard in a non-compact setting. Here, we propose a general method to construct explicit trajectories which explore the space state with positive probability and which permit to prove a petite-set condition for the compact sets of the state space. An application to an age-structured growth-fragmentation process modelling bacterial growth is also shown.

Keywords: Exponential ergodicity, Harris' Theorem, Doeblin minoration, Non-conservative semigroups, Age-size structured equation, Degenerate PDMP.

MSC2020 subject classifications: 60J25, 45C05, 45K05, 92D25.

1 Introduction

The need to include age as a structuring variable in the description of population dynamics has come to be a useful strategy for modellers searching to account for non-Markovian behaviours in a Markovian setting. In particular, in the context of biological applications, the arising of high-throughput single-cell techniques has allowed microbiologists to follow heterogenous populations of isolated bacteria (where the structure is given by their length, biological markers or any other observable) through time. Thereby, this also grants access to the age structure and has put in evidence the non-trivial dependence of age and the other observables at the individual and population scales. The most recent models of bacterial growing include then some sort of age variable, which might not correspond exactly with the *chronological age*, but which might rather evolve in time as a function of the individual trait. This *age* variable still

*CMAP, CNRS, École polytechnique, Institut Polytechnique de Paris, 91120 Palaiseau, France.
E-mail: ignacio.madrid-canales@polytechnique.edu

obeys a renewal equation, which justifies nonetheless its name. This is why we need to extend stability results which account for the level of generality imposed by those models.

In this spirit, we study the long-time behaviour of a stochastic process modelling non-conservative population dynamics which are formalised as a measure-valued process $(Z_t)_{t \geq 0}$ with values in the point measures over \mathbb{R}_+^2 , $\mathcal{M}_p(\mathbb{R}_+^2)$, which represents the age and size of the individuals. For every instant $t > 0$ we can write

$$Z_t = \sum_{i \leq \langle Z_t, 1 \rangle} \delta_{\mathbf{x}_i}, \quad (1.1)$$

where $\mathbf{x}_i = (a_i, y_i)$ denotes the vector trait of individual i , consisting in its age a_i and size y_i . We suppose that for every $f \in C_b^{1,1}(\mathbb{R}_+^2)$, Z_t decomposes as a semi-martingale of the form

$$\langle Z_t, f \rangle \stackrel{\text{def}}{=} \int_{\mathbb{R}_+^2} f(\mathbf{x}) Z_t(d\mathbf{x}) = \langle Z_0, f \rangle + \int_0^t \langle Z_s, \mathcal{Q}f \rangle ds + \mathcal{M}_t^f, \quad (1.2)$$

where \mathcal{M}_t^f is a squared-integrable martingale, and \mathcal{Q} is given for every $f \in C_b^{1,1}(\mathbb{R}_+^2)$ by

$$\mathcal{Q}f(\mathbf{x}) = g(\mathbf{x})^\top \nabla f(\mathbf{x}) + \beta(\mathbf{x}) \left(\int_0^\infty f(0, z) k(\mathbf{x}, z) dz - f(\mathbf{x}) \right) \quad \forall \mathbf{x} \in \mathbb{R}_+^2 \quad (1.3)$$

The first term of the generator \mathcal{Q} is a deterministic transport term, where $g : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ represents a deterministic individual cell growth rate, so that between jumps the variable \mathbf{x} evolves as $\mathbf{x}'(t) = g(\mathbf{x}(t))$. The second term represents the stochastic reproduction events where $\beta : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ is the reproduction rate. The function $k : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ is a positive integrable function modelling the stochastic replacement of the size (by fragmentation or mutation) of individuals of trait \mathbf{x} . The age variable, on the other hand, resets at 0 at each jump. This means that the transition kernel over \mathbb{R}_+^2 is degenerate, of form $\mathbf{x} \mapsto \delta_0(da) \otimes k(\mathbf{x}, z) dz$.

Our goal is to obtain the long-time behaviour of the first-moment semigroup $M_t f(\mathbf{x}) := \mathbb{E}_{\delta_{\mathbf{x}}} [\langle Z_t, f \rangle]$. In particular, we prove a **Malthusian behaviour**:

$$M_t f(\mathbf{x}) = h(\mathbf{x}) e^{\lambda t} \langle \pi, f \rangle + O\left(e^{(\lambda - \omega)t}\right), \quad (1.4)$$

which shows the convergence of $e^{-\lambda t} M_t$ towards a unique stationary measure π at an exponential rate. The parameter $\lambda > 0$ is called the Malthus parameter and represents the growth rate of the population, so that $e^{-\lambda t}$ allows to rescale the mean population size as $t \rightarrow +\infty$. The function h propagates the effect of the initial structure of the population. The constant ω indicates the velocity of the exponential convergence towards π .

Different methods have been developed during the recent years to prove this behaviour: spectral methods, as reviewed in [21] (see for example [23] for an application to a close model); others based on the study of the associated semigroup by Harris' theorem as proposed in some general frameworks by [3, 5, 2] with recent applications in the models considered by [4, 25, 7]. We will follow the latter methods, using the criteria established by Meyn and Tweedie [20], namely: a petite-set condition (Section 2.2) and the existence of a Lyapunov function (Section 2.3). This methods present an alternative to PDE techniques, where criteria based on the probabilistic control of moments replace the harder to obtain Poincaré-type inequalities.

We explore two directions left open in the previous applications, which represent also the sources of our major technical issues: first, the bi-dimensionality of the dynamics, and second, the degeneracy of the transition kernel. Indeed, the underlying stochastic process consists on unidimensional trajectories over a two-dimensional space. Hence, to

uniformly bound in probability the region explored by these trajectories with respect to a non-degenerate measure is not trivial. Similar difficulties have been found for other two-dimensional models such as [13, 26, 8]. Here, we propose to construct explicit trajectories and to average them in time with respect to a nice sampling measure. The inclusion of time permits to compensate the lack of stochasticity of the degenerate jump-transport dynamics with the stochasticity of the reproduction times. More exactly, the utilisation of a petite-set condition instead of a small-sets one is key to obtain the convergence in this two-dimensional setting.

Moreover, compared to the previous works mentioned above, the probabilistic framework brings naturally to work with the operator \mathcal{Q} instead of its dual, as in the more classical PDE settings. Thus, this work lies also in the framework of measure solutions as rigorously developed for example in [14] for the one-dimensional conservative case. Moreover, only the existence of eigenelements for \mathcal{Q} is needed to be able to compute the Doob h -transform and use Harris' theorem. Then, the existence of the *direct* eigenfunction associated to the classical PDE is a consequence of our main result. Our method is then in the spirit of [5], where the authors could exploit known results of existence of the dual eigenelements in the one-dimensional case provided by [1, 10]. In our case, we will have to adapt the latter to the two-dimensional degenerate case studied here.

In particular, we will apply our method to determine the exponential convergence towards a stable size distribution in a bacterial proliferation model called the *adder* model [24, 19, 16, 15]. Individual cells are structured by their added size a which renews to 0 at each division, and their size y which evolves deterministically at exponential rate. The existence of a steady-state distribution and its form was already known since [16] and was recently justified using entropy methods by [15]. Since the eigenelements of the generator are known in this case, by the direct application of Harris' theorem, our method permits to evade technical issues linked to the lack of compactness of the model, which make a classic treatment by PDE and hypocoercivity methods harder to prove and less general.

2 Malthusian behaviour

We are interested in the average dynamics as given by first-moment semigroup M_t defined for every test function $f \in C_b^{1,1}(\mathbb{R}_+^2)$ by:

$$M_t f(\mathbf{x}) := \mathbb{E}[\langle Z_t, f \rangle | Z_0 = \delta_{\mathbf{x}}] \quad \forall \mathbf{x} \in \mathbb{R}_+^2 \quad (2.1)$$

Using Markov's property it's easy to see that M_t verifies the semigroup property. However it is not a Markovian semigroup since it does not necessarily preserve mass (we say it is non conservative). Moreover, using the semi-martingale decomposition 1.2, we verify that M_t is the semigroup associated to the extended generator \mathcal{Q} . This is, for every test function $f \in C^{1,1}(\mathcal{X})$, it is the weak solution of Kolmogorov's equations

$$\partial_t M_t f = M_t \mathcal{Q} f = \mathcal{Q} M_t f. \quad (2.2)$$

Moreover, for any finite measure μ we define the dual semigroup as the measure νM_t given by:

$$(\nu M_t) f := \nu(M_t f) = \int_{\mathcal{X}} M_t f(\mathbf{x}) \nu(d\mathbf{x})$$

So by definition we have $(\mu M_t) f = \mu(M_t f)$ which we write as $\mu M_t f$.

Our main result states the Malthusian behaviour of the semigroup by means of the classical version of Harris theorem as stated in Theorem 6.1 of [20], which we recall below in Theorem 2.1.

Theorem 2.1 (Harris' Theorem (Theorem 6.1 of [20])). *Let $(X_t)_t$ be a right-continuous Markov process with values in some locally compact separable metric space E equipped with its Borelian set $\mathcal{B}(E)$, and let \mathcal{A} be the infinitesimal generator of X . We call P_t the associated transition semigroup. If the two following conditions are verified:*

(H1) Doeblin minoration condition for all compacts.

All compact sets of E are petite for a skeleton chain of X . This is, for every compact set $\mathcal{K} \subset E$ there's a probability mass distribution $\mu = (\mu_n)_{n \in \mathbb{N}}$ over \mathbb{N} and some $\Delta > 0$ such that there exists a non-trivial measure ν (which might depend on Δ and μ) over $\mathcal{B}(E)$ that for every $\mathbf{x} \in \mathcal{K}$ gives the following lower bound:

$$\langle \mu, \delta_{\mathbf{x}} P_{\Delta} f \rangle = \sum_{n \in \mathbb{N}} \mu_n P_{n\Delta} f(\mathbf{x}) \geq \langle \nu, f \rangle.$$

(H2) Foster-Lyapunov drift condition.

There exist a coercive function V and some $c > 0, d < \infty$ such that

$$\mathcal{A}V(\mathbf{x}) \leq -cV(\mathbf{x}) + d \quad \forall \mathbf{x} \in E,$$

Then, there exist a unique non-trivial probability measure π and $C, \omega > 0$ such that for every $\mathbf{x} \in E$ and $t \geq 0$

$$\|\delta_{\mathbf{x}} P_t - \pi\|_{1+V} \leq C(1 + V(\mathbf{x})) \exp(-\omega t) \quad (2.3)$$

Remark that we need a Markovian (conservative) semigroup. To overcome this problem, similarly as in [5], we perform a so-called Doob h -transform, to obtain a conservative semigroup P_t with the dynamics of M_t . To do so, we require first to have some pair (λ, h) such that $\mathcal{Q}h = \lambda h$ and $h > 0$. Then, using such pair we define

$$P_t f(\mathbf{x}) := \frac{M_t(hf)(\mathbf{x})}{e^{\lambda t} h(\mathbf{x})}. \quad (2.4)$$

Then we can come back the ergodic behaviour of $(M_t)_{t \geq 0}$ by looking at the limit of $M_t f = e^{\lambda t} h P_t (f/h)$. In particular, the generator associated with P_t is given explicitly by Eq. (2.5).

Proposition 2.2. *Suppose the existence of a pair (λ, h) , $\lambda > 0, h > 0$ such that $\mathcal{Q}h = \lambda h$. Then, P_t defined by Eq. (2.4) is a positive Markovian semigroup whose infinitesimal generator is given for $f \in C_b^{1,1}(\mathbb{R}_+^2)$ by*

$$\mathcal{A}f(\mathbf{x}) = g(\mathbf{x})^\top \nabla f(\mathbf{x}) + \beta(\mathbf{x}) \left(\int_0^\infty [f(0, z) - f(\mathbf{x})] \frac{h(0, z)}{h(\mathbf{x})} k(\mathbf{x}, z) dz \right) \quad \forall \mathbf{x} \in \mathbb{R}_+^2 \quad (2.5)$$

Proof. By definition and evaluating at $t = 0$ we have:

$$\begin{aligned} \mathcal{A}f(\mathbf{x}) &= \left. \frac{\partial}{\partial t} P_t f \right|_{t=0}(\mathbf{x}) \\ &= \left. \frac{M_t \mathcal{Q}(hf)}{e^{\lambda t} h} - \frac{\lambda M_t(hf)}{e^{\lambda t} h} \right|_{t=0}(\mathbf{x}) \\ &= \frac{\mathcal{Q}(hf)(\mathbf{x})}{h(\mathbf{x})} - \lambda f(\mathbf{x}) \end{aligned}$$

Then, using the value of \mathcal{Q} applied to hf and that $\mathcal{Q}h = \lambda h$ we get

$$\begin{aligned}
\frac{\mathcal{Q}(hf)(\mathbf{x})}{h(\mathbf{x})} &= \frac{f(\mathbf{x})}{h(\mathbf{x})} g(\mathbf{x})^\top \nabla h(\mathbf{x}) + g(\mathbf{x})^\top \nabla f(\mathbf{x}) + \beta(\mathbf{x}) \left(\int_0^\infty h(0, z) f(0, z) \frac{k(\mathbf{x}, z)}{h(\mathbf{x})} dz - f(\mathbf{x}) \right) \\
&= \frac{f(\mathbf{x})}{h(\mathbf{x})} (g(\mathbf{x})^\top \nabla h(\mathbf{x}) - \beta(\mathbf{x}) h(\mathbf{x})) + g(\mathbf{x})^\top \nabla f(\mathbf{x}) + \beta(\mathbf{x}) \int_0^\infty h(0, z) f(0, z) \frac{k(\mathbf{x}, z)}{h(\mathbf{x})} dz \\
&= \frac{f(\mathbf{x})}{h(\mathbf{x})} \left(\lambda h(\mathbf{x}) - \beta(\mathbf{x}) \int_0^\infty h(0, z) k(\mathbf{x}, z) \right) + g(\mathbf{x})^\top \nabla f(\mathbf{x}) \\
&\quad + \beta(\mathbf{x}) \int_0^\infty h(0, z) f(0, z) \frac{k(\mathbf{x}, z)}{h(\mathbf{x})} dz \\
&= \lambda f(\mathbf{x}) + g(\mathbf{x})^\top \nabla f(\mathbf{x}) + \beta(\mathbf{x}) \int_0^\infty [f(0, z) - f(\mathbf{x})] \frac{h(0, z)}{h(\mathbf{x})} k(\mathbf{x}, z) dz
\end{aligned}$$

Finally, subtracting $\lambda f(\mathbf{x})$ we obtain the form of generator \mathcal{A} . \square

Hence, the work is structured as follows: first, in Section 3 we prove the existence of a pair (λ, h) which solves the eigenvalue problem $\mathcal{Q}h = \lambda h$ under the assumptions 3.3. The same set of Assumptions allows us to prove the Doeblin condition (H1) in Section 4. We do not provide a general Foster-Lyapunov condition (H2), suitable for our general case. However, we show its existence in our application to a growth-fragmentation model in Section 5. This last model has already been studied since the works of [16], and the exponential convergence has been recently shown in [15] using Generalized Relative Entropy techniques. Here, we show that the knowledge of the eigenelements (λ, h) for the generator allows to provide a simpler proof of convergence using Harris' theorem. Indeed, the arguments presented in Section 3 can be avoided when the existence of eigenelements is known apriori, which might be the case in several practical applications. Nonetheless, our general method allows us to give an answer to one of the perspectives listed by [15], who couldn't generalise their argument in the case of a general drift function g . Thus, our main result reads as follows:

Theorem 2.3 (Exponential ergodicity). *Under Assumptions 3.3 and if the Lyapunov-Foster condition (H2) of Theorem 2.1 is verified for some coercive function $V : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$, there is a unique probability measure π such that there exist constants $C, \omega, \Lambda > 0$ which verify for every initial condition $\mu_0 \in \mathcal{M}_p(\mathbb{R}_+^2)$*

$$\|e^{-\Lambda t} \mu_0 M_t - \langle \mu_0, h \rangle \pi\|_{1+V} \leq C(1 + \mu_0 V) e^{-\omega t} \quad (2.6)$$

where $\|\cdot\|_V$ is a weighted total variation norm defined by

$$\|u\|_V = \int_0^\infty \int_0^y V(a, y) |u(a, y)| da dy.$$

Moreover, π is absolutely continuous with respect to the Lebesgue measure.

3 Preliminary definitions and assumptions

We begin by recalling some useful properties of the deterministic flow, which are classical results for an autonomous system of first order ODE (refer for example to Theorem D.1 of [18]):

Lemma 3.1 (Flow properties and notations.). *Let $\mathbf{x} \in \mathbb{R}_+^2$. Consider $g = (g_1, g_2) \in C^1(\mathbb{R}_+^2)$ and suppose that $g_1 > 0$. The autonomous first-order system of Ordinary Differential Equations (ODE)*

$$\begin{aligned}
\frac{d\mathbf{u}(t)}{dt} &= g(\mathbf{u}(t)), \quad t \in \mathbb{R} \\
\mathbf{u}(0) &= \mathbf{x}
\end{aligned} \quad (3.1)$$

defines a unique flow $\varphi^t : \mathcal{X} \ni \mathbf{x} \mapsto \varphi^t(\mathbf{x}) \in \mathcal{X}$ which is the solution $\mathbf{u}(t)$ of (3.1) at time t with initial condition $\mathbf{x} \in \mathcal{X}$ where $\mathcal{X} = \bigcup_{y \geq 0} \Gamma_{(0,y)}^+$ where $\Gamma_{\mathbf{x}}^+$ will be defined below. We write $\varphi^t = (\varphi_1^t, \varphi_2^t)$ for the marginal flows of the age and size. We define then $\Gamma_{\mathbf{x}}^+ = \{\varphi^t(\mathbf{x}), t \geq 0\}$ and $\Gamma_{\mathbf{x}}^- = \{\varphi^t(\mathbf{x}), t \leq 0\}$ and call $\Gamma_{\mathbf{x}} = \Gamma_{\mathbf{x}}^+ \cup \Gamma_{\mathbf{x}}^-$ the unique orbit passing through \mathbf{x} . Moreover:

1. The flow is a group in the time variable: $\varphi^t \varphi^s = \varphi^{t+s} = \varphi^s \varphi^t$, $\varphi^0 = \text{Id}$, and has inverse $(\varphi^t)^{-1} = \varphi^{-t}$, which is the solution to the ODE $\mathbf{u}'(t) = -g(\mathbf{u}(t))$.
2. The flow depends smoothly on the initial conditions: $\forall t \in \mathbb{R}$, $\varphi^t \in C^1(\mathbb{R}_+^2)$. In particular the Jacobian matrix of the flow with respect to the initial condition is given explicitly by

$$\mathcal{D}\varphi^t(\mathbf{x}) = \exp\left(\int_0^t \mathcal{D}g(\varphi^s(\mathbf{x})) ds\right).$$

where \mathcal{D} denotes the Jacobian matrix with respect to $\mathbf{x} = (a, y)$, and $\exp(\cdot)$ corresponds here to a matrix exponential.

3. For all fixed $\mathbf{x} = (a_0, y_0) \in \mathcal{X}$, if $g_1 > 0$, then there is a unique function $Y_{\mathbf{x}} : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for all $(a, y) \in \Gamma_{\mathbf{x}}$, we have $Y_{\mathbf{x}}(a) = y$. This represents the size at a given age of an individual with initial condition \mathbf{x} . In other words, for all $t \geq 0$,

$$\varphi^t(\mathbf{x}) = (a(t), Y_{\mathbf{x}}(a(t))).$$

Moreover, $Y_{\mathbf{x}} \in C^1(\mathbb{R}_+)$ and it is solution of the first order one-dimensional ODE

$$Y_{\mathbf{x}}'(a) = \frac{g_2(a, Y_{\mathbf{x}}(a))}{g_1(a, Y_{\mathbf{x}}(a))} \quad ; \quad Y_{\mathbf{x}}(a_0) = y_0$$

Analogously, one defines its inverse function $A_{\mathbf{x}}(y)$ which gives the age at size y for an individual with initial condition \mathbf{x} , and hence verifies

$$\varphi^t(\mathbf{x}) = (A_{\mathbf{x}}(y(t)), y(t)), \quad t \geq 0.$$

4. For all fixed $\mathbf{x} \in \mathcal{X}$, we write $\phi_{\mathbf{x}}(t) := \varphi^t(\mathbf{x})$ as a function of time (from \mathbb{R} to \mathbb{R}_+^2). Then, the inverse function $\phi_{\mathbf{x}}^{-1} : \Gamma_{\mathbf{x}} \rightarrow \mathbb{R}$ such that $\phi_{\mathbf{x}}^{-1}(\phi_{\mathbf{x}}(t)) = t$ is well defined. For every $\mathbf{x}_0 \in \mathcal{X}$ and $\mathbf{x}_1 \in \Gamma_{\mathbf{x}_0}$ we read $\phi_{\mathbf{x}_0}^{-1}(\mathbf{x}_1)$ as the time needed along $\Gamma_{\mathbf{x}_0}$ to go from \mathbf{x}_0 to \mathbf{x}_1 . Moreover if we write $\mathbf{x}_0 = (a_0, y_0)$, $\mathbf{x}_1 = (a_1, y_1)$, this quantity is given by

$$\phi_{\mathbf{x}_0}^{-1}(\mathbf{x}_1) = \int_{a_0}^{a_1} \frac{1}{g_1(a, Y_{\mathbf{x}_0}(a))} da = \int_{y_0}^{y_1} \frac{1}{g_2(A_{\mathbf{x}_0}(y), y)} dy$$

Importantly, for the set of assumptions given below, we have $0 < \phi_{\mathbf{x}_0}^{-1}(\mathbf{x}_1) < \infty$ for all $\mathbf{x}_0 \in \mathcal{X} \setminus \{0\}$ and $\mathbf{x}_1 \in \Gamma_{\mathbf{x}_0}$.

Let us also consider the following probability space which will be useful to compute and interpret some of the estimates which will be obtained below.

Definition 3.2. Consider a probability space $(\mathbb{R}_+, \mathcal{B}(\mathbb{R}_+), \mathbb{P}_{\mathbf{x}})$ in which the random couple $(T, Z) \in \mathbb{R}_+ \times \mathbb{R}_+$ gives the first jump time T and size Z after this first jump of a trajectory beginning at $\mathbf{x} \in \mathcal{X}$. Hence, for all $\mathbf{x} \in \mathcal{X}$, the couple (T, Z) has joint probability density

$$p_{\mathbf{x}}(t, z) = \frac{1}{C_{\mathbf{x}}} k(\varphi^t(\mathbf{x}), z) \psi(t|\mathbf{x}),$$

where the normalisation constant is given by

$$C_{\mathbf{x}} = \int_0^{\infty} \int_0^{\infty} k(\varphi^t(\mathbf{x}), z') \psi(t|\mathbf{x}) dt dz',$$

which is the mean number of offspring produced by an individual of initial configuration \mathbf{x} after its first jump, and

$$\psi(t|\mathbf{x}) = \beta(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds\right)$$

is the marginal probability density of the time of the first jump, conditionally to the initial configuration \mathbf{x} , and which is well defined for the set of assumptions given below. We write $\mathbb{E}_{\mathbf{x}}$ the associated expectation. Fig. 1 summarises the definitions introduced in this section.




Figure 1: Flow notations introduced in Lemma 3.1 and the probabilistic definition of the random couple (T, Z) introduced in Definition 3.2.

Now, let us consider the following set of assumptions, whose biological meaning and implications are commented below.

Assumptions 3.3. Assume that we have

- (i) *Smooth and uniformly controlled flow:* $g = (g_1, g_2) \in C^1(\mathbb{R}_+^2)$, $g_1 > 0$ and there are some constants $c_0, c_1, c_2 > 0$ such that for all $(a, y) \in \mathbb{R}_+^2$

$$g_1(a, y) \geq c_0 a, \quad g_1(a, y) \leq c_1(1 + a), \quad g_2(a, y) \leq c_1(1 + y),$$

$$|\partial_a g_i(a, y)| \leq c_2(1 + a + y), \quad |\partial_y g_i(a, y)| \leq c_2(1 + a + y).$$

and for all $y > 0, a \geq 0$, we have $g_2(a, y) \leq g_2(0, y)$.

- (ii) *Regular reproduction rate:* $\beta \in C(\mathbb{R}_+^2, \mathbb{R}_+)$, and $B = \beta/g_1 \in C(\mathbb{R}_+^2)$, such that there are constants $a^*, \beta_-, \beta_+ > 0$ s.t. for all $a > a^*y \geq 0, \beta_- < B(a, y) < \beta_+$, and $B(a, y) = 0$ for all $a \leq a^*$.

- (iii) *Regular transition kernel:* For all $z \geq 0, \mathbf{x} \mapsto k(\mathbf{x}, z)$ is a continuous function on \mathbb{R}_+^2 , and for all $\mathbf{x} \in \mathbb{R}_+^2, z \mapsto k(\mathbf{x}, z)$ is a continuous function on \mathbb{R}_+ . The total offspring of individuals of trait \mathbf{x} is $\|k(\mathbf{x}, \cdot)\|_1 := \int_0^{+\infty} k(\mathbf{x}, z) dz$ with $1 < \|k(\mathbf{x}, \cdot)\|_1 \leq \bar{K}$ for all $\mathbf{x} \in \mathbb{R}_+^2$. In particular, we consider two distinct cases:

- (a) *Fragmentation kernel:* For all $a \geq 0, \text{supp } k(a, y, \cdot) \subseteq (0, y)$.
- (b) *Compactly supported mutational kernel:* It exists a compact set $S \subset \mathbb{R}_+$ such that for all $a \geq 0, \text{supp } k(a, y, \cdot) \subseteq S$, and some interval $I \subset \mathbb{R}_+$ and $\epsilon_0 > 0$ such that for all $y \in S$ and $z \in S \cap \mathbb{B}_{\epsilon_0}(y)$, the open ball of radius ϵ_0 around y , we have $I \subset \{a > 0 : \beta(a, y)k(a, y, z) > 0\}$.

(iv) *Lower bounded transition kernel:* For all fixed value of $z > 0$, there exists some non-empty open interval $D(z)$ with length bounded between δ_- and δ_+ , both independent of z , and a positive value $\varepsilon(z)$ such that for all $\mathbf{x} \in \mathbb{R}_+^2$, $k(\mathbf{x}, z) > \varepsilon(z)\mathbb{1}_{D(z)}(\mathbf{x})$.

We comment on the meaning of these assumptions. Assumption 3.3-(i) ensures that the size and age do not explode in finite time. The control on the derivatives will also allow to control the influence of the initial conditions on the flow (Lemma 3.1.2). Assumption 3.3-(ii) allows to write the division rate as $\beta(\mathbf{x}) = g_1(\mathbf{x})B(\mathbf{x})$ where function B should be interpreted as an "age hazard rate". Indeed, if we let \mathbf{A} be the random variable representing the age at division of an individual, then under $\mathbb{P}_{\mathbf{x}}$ a change of variables gives

$$\mathbb{P}_{\mathbf{x}}(\mathbf{A} \geq a) = \exp\left(-\int_0^a B(\alpha, Y_{\mathbf{x}}(\alpha))d\alpha\right). \quad (3.2)$$

Hence for every initial condition $\mathbf{x} \in \mathbb{R}_+^2$ we have

$$B(a, Y_{\mathbf{x}}(a)) da = \mathbb{P}_{\mathbf{x}}(\mathbf{A} \in [a, a + da] | \mathbf{A} \geq a).$$

Proof of Eq. (3.2). Let us recall that under $\mathbb{P}_{\mathbf{x}}$ introduced in Definition 3.2, the random variable T is the first jump time. Then, since $\phi_{\mathbf{x}}^{-1}(a, Y_{\mathbf{x}}(a))$ is the time needed to reach age a starting from an initial configuration \mathbf{x} we have

$$\begin{aligned} \mathbb{P}_{\mathbf{x}}(\mathbf{A} \geq a) &= \mathbb{P}_{\mathbf{x}}(T \geq \phi_{\mathbf{x}}^{-1}(a, Y_{\mathbf{x}}(a))) \\ &= \exp\left(-\int_0^{\phi_{\mathbf{x}}^{-1}(a, Y_{\mathbf{x}}(a))} \beta(\varphi^s(\mathbf{x})) ds\right) \end{aligned}$$

Then, doing the change of variables $s \mapsto \alpha(s) = \varphi_1^s(\mathbf{x})$, i.e. changing from time s , to the age $\alpha(s)$ at time s starting from \mathbf{x} , and since $\alpha'(s) = \frac{d}{ds}\varphi_1^s(\mathbf{x}) = g_1(\varphi_1^s(\mathbf{x}))$ we have

$$\begin{aligned} \int_0^{\phi_{\mathbf{x}}^{-1}(a, Y_{\mathbf{x}}(a))} \beta(\varphi^s(\mathbf{x})) ds &= \int_0^a \beta(\alpha, Y_{\mathbf{x}}(\alpha)) \frac{1}{g_1(\alpha, Y_{\mathbf{x}}(\alpha))} d\alpha \\ &= \int_0^a B(\alpha, Y_{\mathbf{x}}(\alpha)) d\alpha \end{aligned}$$

and hence

$$\mathbb{P}_{\mathbf{x}}(\mathbf{A} \geq a) = \exp\left(-\int_0^a B(\alpha, Y_{\mathbf{x}}(\alpha))d\alpha\right).$$

□

Biologically this has been interpreted as individuals not perceiving actual time, but rather their own biological age, upon which the division event is decided [24]. The parameter a^* is the minimal division age. It imposes that it is not possible to divide immediately after birth. For ages bigger than a^* , the bounds on B allow to stochastically bound the age at division between two exponential random variables of rate parameter β_- and β_+ . Assumption 3.3-(iii) imposes inexact, not perfectly mitotic cell divisions which give always two new individuals. The two considered cases bring together a broad family of transition kernels used in similar models. In particular the assumptions concerning the mutational kernel are inspired from [23]. Importantly, the compactness is needed to prove the existence of the eigenelements of \mathcal{Q} but not for the Doeblin minoration, which holds in more general cases. In this line, Assumption 3.3-(iv) is key to obtain the Doeblin minoration condition and generalises similar requirements needed in the one-dimensional case, such as Eq. (8) of [5] for auto-similar fragmentation kernels, and Eq. (10) of [6], or Assumption (A4) of [23] for general non-local mutation-type kernels.

4 Existence of the eigenelements of \mathcal{Q}

Now, in order to bring ourselves to the conservative setting, we begin by showing the existence of some pair of eigenelements for \mathcal{Q} .

Proposition 4.1 (Existence of eigenelements). *Under Assumptions 3.3, there exist a positive constant $\lambda > 0$ and a positive function $h \in W_{loc}^{1,\infty}(\mathcal{X})$ such that*

$$\mathcal{Q}h = \lambda h.$$

To do so, we can reformulate the eigenvalue problem as a one-dimensional fixed point problem. This is a classical strategy and other applications in two-dimensional spaces can be found for example in [9, 17, 15]. In particular, we follow closely the arguments of [9] which corresponds to the case $g_1 \equiv 1$ with a fragmentation kernel and with additional confinement assumption in the drift term which would allow us to work in a compact interval in one of the two dimensions. We generalise this approach here.

Lemma 4.2 (Reformulation as a renewal equation). *Any pair (λ, h) such that $\lambda > 0$ and $h \in W_{loc}^{1,\infty}(\mathcal{X})$ is solution almost everywhere to $\mathcal{Q}h = \lambda h$ and verifies*

$$\lim_{t \rightarrow +\infty} h(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) = 0 \quad (4.1)$$

if and only if it verifies the renewal formula

$$h(\mathbf{x}) = \int_0^\infty h(0, z) K_\lambda(\mathbf{x}, z) dz, \quad (4.2)$$

where

$$K_\lambda(\mathbf{x}, z) = C_{\mathbf{x}} \int_0^\infty e^{-\lambda t} p_{\mathbf{x}}(t, z) dt \quad (4.3)$$

Remark 4.3. Using Definition 3.2 we can then write Eq. (4.2) as

$$h(\mathbf{x}) = C_{\mathbf{x}} \mathbb{E}_{\mathbf{x}}[h(0, Z)e^{-\lambda T}]. \quad (4.4)$$

Proof of Lemma 4.2. We proceed by the method of integration along characteristics. Since $h \in W_{loc}^{1,\infty}$, taking weak derivatives we obtain that, almost everywhere:

$$\begin{aligned} & \frac{\partial}{\partial t} \left(h(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) \right) \\ &= (\nabla h(\varphi^t(\mathbf{x}))^\top g(\varphi^t(\mathbf{x})) - (\beta(\varphi^t(\mathbf{x})) + \lambda)h(\varphi^t(\mathbf{x}))) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) \\ &= \left(\mathcal{Q}h(\varphi^t(\mathbf{x})) - \lambda h(\varphi^t(\mathbf{x})) - \beta(\varphi^t(\mathbf{x})) \int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \right) e^{-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t} \quad (4.5) \end{aligned}$$

First, suppose that (λ, h) is solution a.e. to $\mathcal{Q}h = \lambda h$ and that Eq. (4.1) is verified. Then, integrating Eq. (4.5) in $(0, +\infty)$ and using the decay condition Eq. (4.1) results into

$$h(\mathbf{x}) = \int_0^{+\infty} \beta(\varphi^t(\mathbf{x})) \int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) dt$$

which, by Fubini, gives exactly Eq. (4.2). Now, suppose that we have Eq. (4.2). Then we

have:

$$\begin{aligned}
h(\varphi^t(\mathbf{x})) &= \int_0^\infty h(0, z) K_\lambda(\varphi^t(\mathbf{x}), z) dz \\
&= \int_0^\infty \int_0^\infty h(0, z) k(\varphi^{t+s}(\mathbf{x}), z) \beta(\varphi^{t+s}(\mathbf{x})) \exp\left(-\int_t^{t+s} \beta(\varphi^u(\mathbf{x})) du - \lambda s\right) ds dz \\
&= \left(\int_0^\infty \int_0^\infty h(0, z) k(\varphi^s(\mathbf{x}), z) \beta(\varphi^s(\mathbf{x})) \exp\left(-\int_0^s \beta(\varphi^u(\mathbf{x})) du - \lambda s\right) ds dz \right. \\
&\quad \left. - \int_0^\infty \int_0^t h(0, z) k(\varphi^s(\mathbf{x}), z) \beta(\varphi^s(\mathbf{x})) \exp\left(-\int_0^s \beta(\varphi^u(\mathbf{x})) du - \lambda s\right) ds dz \right) \\
&\quad \times \exp\left(\int_0^t \beta(\varphi^u(\mathbf{x})) du + \lambda t\right)
\end{aligned}$$

Therefore, using Eq. (4.2):

$$h(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) = h(\mathbf{x}) - \int_0^\infty \int_0^t h(0, z) k(\varphi^s(\mathbf{x}), z) \psi(s|\mathbf{x}) e^{-\lambda s} ds dz \quad (4.6)$$

As $t \rightarrow +\infty$, the improper integral in the RHS of Eq. (4.6) converges towards

$$\lim_{t \rightarrow +\infty} \int_0^\infty \int_0^t h(0, z) k(\varphi^s(\mathbf{x}), z) \psi(s|\mathbf{x}) e^{-\lambda s} ds dz = \int_0^\infty h(0, z) K_\lambda(\mathbf{x}, z) dz = h(\mathbf{x}),$$

from which we obtain Eq. (4.1). Moreover, supposing that $h \in W_{\text{loc}}^{1, \infty}$ and taking weak derivatives in Eq. (4.6) we obtain

$$\frac{\partial}{\partial t} \left(h(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda t\right) \right) = - \int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) \psi(t|\mathbf{x}) e^{-\lambda t} dz$$

Hence, a comparison with Eq. (4.5) gives that for all $\mathbf{x} \in \mathcal{X}$ and $t > 0$ we have almost everywhere $\mathcal{Q}h(\varphi^t(\mathbf{x})) - \lambda h(\varphi^t(\mathbf{x})) = 0$, or equivalently, $\mathcal{Q}h = \lambda h$ almost everywhere in \mathcal{X} . \square

Remark 4.4. In particular the function $\eta(y) := h(0, y)$ defined for all $y \geq 0$ is solution to the fixed point problem

$$\eta(y) = \int_0^\infty \eta(z) K_\lambda(0, y, z) dz. \quad (4.7)$$

Therefore we will consider the operator G_λ defined for $f \in C^1(\mathbb{R}_+)$ by

$$\mathcal{G}_\lambda f(y) = \int_0^\infty f(z) K_\lambda(0, y, z) dz \quad \forall y > 0. \quad (4.8)$$

We also introduce the operator $\mathcal{J}_\lambda : \mathcal{M}(\mathbb{R}_+) \rightarrow \mathcal{M}(\mathbb{R}_+)$ which for any Radon measure ν supported in \mathbb{R}_+ gives

$$\mathcal{J}_\lambda \nu = \left(\int_0^\infty K_\lambda(0, z, y) \nu(dz) \right) dy \quad (4.9)$$

and verifies the duality property below:

Proposition 4.5. For every $\lambda \geq 0$, \mathcal{J}_λ is the adjoint operator of \mathcal{G}_λ .

Proof. Let $f \in C(\mathbb{R}_+^2)$ and $\nu \in \mathcal{M}(\mathbb{R}_+^2)$. By Fubini's Theorem,

$$\begin{aligned}
\langle \nu, \mathcal{G}_\lambda f \rangle &= \int_0^\infty \left(\int_0^\infty f(z) K_\lambda(0, y, z) dz \right) \nu(dy) \\
&= \int_0^\infty f(z) \left(\int_0^\infty K_\lambda(0, y, z) \nu(dy) \right) dz = \langle \mathcal{J}_\lambda \nu, f \rangle.
\end{aligned}$$

\square

Remark 4.6. From Eq. (4.4), we can write

$$\mathcal{G}_\lambda f(y) = C_{(0,y)} \mathbb{E}_{(0,y)}[f(Z)e^{-\lambda T}].$$

where again $C_{(0,y)} = \|K_0(0,y, \cdot)\|_1$ is the mean number of offspring produced by an individual of initial size y after its first jump.

Proof of Proposition 4.1. We aim to prove that there is a unique $\lambda > 0$ for which the operator \mathcal{G}_λ admits a unique fixed point $h(0, \cdot)$. The pair (λ, h) is then solution to the eigenproblem $\mathcal{Q}h = \lambda h$. This will be proven by means of Krein-Rutman's theorem. In order to be able to apply this theorem we need to work with a strictly positive compact operator. For the compactly supported mutational kernel it is immediately the case, however it is not the case for \mathcal{G}_λ with a fragmentation kernel. Thus, we shall follow a standard approximation scheme for the proof which is structured as follows:

1. We define a truncated version of \mathcal{G}_λ which by Arzela-Ascoli's theorem we prove to be a positive compact operator in the Banach space of continuous functions.
2. We apply Krein-Rutman theorem to prove that for each $\lambda \geq 0$ the truncated operator admits a unique eigenvalue $\mu_\lambda \geq 0$ and suitably normalised eigenfunction $h_\lambda \geq 0$.
3. We prove that there exists a unique $\lambda_0 > 0$ such that $\mu_{\lambda_0} = 1$
4. We prove that the value of λ_0 is uniformly bounded for all the members of the family of truncated operators.
5. We pass to the limit and show that the limit eigenelements $(\lambda_0, h_{\lambda_0})$ of the family of truncated operators are indeed solution to the fixed point problem.

Note that the proof is also valid for the compact mutational kernel which verifies Assumption 3.3-(iii)-(b), but in that case neither the truncation nor the uniform estimates are needed.

Step 1 : Construction of the truncated operator.

For each $R > 0$ let $\mathcal{G}_\lambda^R : C^1([0, R]) \rightarrow C^1([0, R])$ defined for all $\lambda > 0$, for $f \in C^1([0, R])$ by

$$\mathcal{G}_\lambda^R f(y) = \int_0^R f(z) K_\lambda^R(0, y, z) dz \quad \forall y \in (0, R) \quad (4.10)$$

with

$$K_\lambda^R(0, y, z) = \int_0^\infty \left(k(\varphi^t(0, y), z) + \frac{1}{R} \int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta \right) \psi(t|(0, y)) e^{-\lambda t} dt$$

We require to add the uniform correction $z \mapsto \frac{1}{R} \int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta$ in order to endorse the strict positivity of the operator. Indeed, for all $y \in [0, R]$, from Fubini's theorem, Assumption 3.3-(iii) and Jensen's inequality we obtain

$$\begin{aligned} \int_0^R K_\lambda^R(0, y, z) dz &= \int_0^\infty \left(\int_0^\infty k(\varphi^t(0, y), z) dz \right) \psi(t|(0, y)) e^{-\lambda t} dt \\ &> \int_0^\infty e^{-\lambda t} \psi(t|(0, y)) dt \\ &\geq \exp(-\lambda \mathbb{E}_{(0,y)}[T]). \end{aligned}$$

Moreover, Assumption 3.3-(ii) gives that

$$0 < \mathbb{E} \left[\phi_{(0,y)}^{-1}(\mathbf{A}_-, Y_{(0,y)}(\mathbf{A}_-)) \right] \leq \mathbb{E}_{(0,y)}[T] \leq \mathbb{E} \left[\phi_{(0,y)}^{-1}(\mathbf{A}_+, Y_{(0,y)}(\mathbf{A}_+)) \right] < +\infty$$

where \mathbf{A}_- (respectively \mathbf{A}_+) follows an Exponential distribution of parameter β_- (respectively β_+). Therefore for all positive $f \in C^1([0, R])$, $G_\lambda^R f > 0$.

Moreover, in analogy with Definition 3.2, we define for all $R > 0$ the random couple $(T_R, Z_R) \in \mathbb{R}_+ \times [0, R]$ such that under $\mathbb{P}_{(0,y)}$ they have joint probability density

$$p_{(0,y)}^R(t, z) = \frac{1}{C_{(0,y)}} \left(k(\varphi^t(0, y), z) + \frac{\mathbb{1}_{z \leq R}}{R} \int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta \right) \psi(t|(0, y)),$$

then we can write

$$\mathcal{G}_\lambda^R f(y) = C_{(0,y)} \mathbb{E}_{(0,y)} [f(Z_R) e^{-\lambda T_R} \mathbb{1}_{Z_R \leq R}]. \quad (4.11)$$

Step 2 : Existence of the eigenelements of \mathcal{G}_λ^R .

We begin by proving that for all $\varepsilon > 0$, $\lambda \geq 0$ and $R > 0$, \mathcal{G}_λ^R is compact. We show that for every sequence $(f_n)_n$ in the unit ball of $C([0, R])$ there exists a subsequence of $(\mathcal{G}_\lambda^R f_n)_n$ which converges in $C([0, R])$ equipped with the uniform norm.

- i. **Uniform bound:** For all $y \in (0, R)$, f in the unit ball of $C[0, R]$ we have from Eq. (4.11):

$$\mathcal{G}_\lambda^R f(y) \leq C_{(0,y)} \|f\|_\infty \leq \bar{K}$$

- ii. **Equicontinuity:** Since $g_1 \in C^1(\mathbb{R}_+^2)$ and is strictly positive, and k is continuous in the first two variables, we have that for every $\lambda \geq 0$, $(y, z) \in [0, R] \times [0, R] \mapsto K_\lambda(0, y, z)$ is an uniformly continuous function on $[0, R] \times [0, R]$. Therefore for all $\lambda \geq 0$ and $\varepsilon > 0$ there exists $\delta > 0$ such that if $|y_1 - y_2| + |z_1 - z_2| < \delta$ for $y_1, y_2, z_1, z_2 \in [0, R]$, then $|K_\lambda(0, y_1, z_1) - K_\lambda(0, y_2, z_2)| < \varepsilon/R$. Hence, for all f in the unit ball, $y_1, y_2 \in [0, R]$ such that $|y_1 - y_2| < \delta$ we have:

$$|\mathcal{G}_\lambda^R f(y_1) - \mathcal{G}_\lambda^R f(y_2)| \leq \int_0^R |f(z)| |K_\lambda(0, y_1, z) - K_\lambda(0, y_2, z)| dz < \varepsilon$$

independently on y_1, y_2 .

Finally, by Ascoli's criterium, there exists a convergent subsequence of $(\mathcal{G}_\lambda^R f_n)_n$ and so the operator \mathcal{G}_λ^R is strictly positive and compact for the uniform topology of $C([0, R])$. Therefore, by Krein-Rutman theorem [22] there exists a unique triplet of a positive real value $\mu_\lambda^R > 0$, function $\eta_\lambda^R > 0$ continuous on $[0, R]$, and a positive Radon measure ν_λ^R supported on $[0, R]$ such that

$$\mathcal{G}_\lambda^R \eta_\lambda^R = \mu_\lambda^R \eta_\lambda^R \quad (4.12)$$

$$\mathcal{J}_\lambda^R \nu_\lambda^R = \mu_\lambda^R \nu_\lambda^R, \nu_\lambda^R([0, R]) = 1 \quad (4.13)$$

$$\langle \nu_\lambda^R, \eta_\lambda^R \rangle_R = 1, \quad (4.14)$$

where we denote $\langle \nu, f \rangle_R = \int_0^R f(y) \nu(dy)$.

Step 3 : Existence and uniqueness of $\lambda_0 > 0$ such that $\mu_{\lambda_0}^R = 1$

We show that the mapping $\lambda \mapsto \mu_\lambda^R$ is a continuous strictly decreasing function which goes through the value of 1 at some point. First, note that from Equations (4.12) and (4.14), we have

$$\langle \nu_\lambda^R, \mathcal{G}_\lambda^R \eta_\lambda^R \rangle_R = \mu_\lambda^R \quad (4.15)$$

We prove that $\lambda \mapsto \langle \nu_\lambda^R, \mathcal{G}_\lambda^R \eta_\lambda^R \rangle_R$ is differentiable continuous and decreasing. Let us consider the derivatives in the sense of distributions $\partial_\lambda \nu_\lambda^R$ and $\partial_\lambda \eta_\lambda^R$. We show below that $\lambda \mapsto \langle \nu_\lambda^R, \mathcal{G}_\lambda^R \eta_\lambda^R \rangle_R$ is actually strongly differentiable with respect to λ as it has the same regularity as $\lambda \mapsto \mathcal{G}_\lambda^R f$. First, by dominated convergence, differentiating under the integral sign on Eq. (4.11) gives for every $f \in C^1([0, R])$,

$$(\partial_\lambda \mathcal{G}_\lambda^R) f(y) = -C_{(0,y)} \mathbb{E}_{(0,y)}[f(Z_R) T_R e^{-\lambda T_R} \mathbb{1}_{Z_R \leq R}]. \quad (4.16)$$

Then, by differentiating under the duality brackets, and using the duality between \mathcal{G} and \mathcal{J} with (4.12) and (4.13), we obtain

$$\begin{aligned} \partial_\lambda \mu_\lambda^R &= \langle \partial_\lambda \nu_\lambda^R, \mathcal{G}_\lambda^R \eta_\lambda^R \rangle + \langle \nu_\lambda^R, \mathcal{G}_\lambda^R (\partial_\lambda \eta_\lambda^R) \rangle + \langle \nu_\lambda^R, (\partial_\lambda \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle \\ &= \langle \partial_\lambda \nu_\lambda^R, \mathcal{G}_\lambda^R \eta_\lambda^R \rangle + \langle \mathcal{J}_\lambda^R \nu_\lambda^R, \partial_\lambda \eta_\lambda^R \rangle + \langle \nu_\lambda^R, (\partial_\lambda \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle \\ &= \mu_\lambda^R (\langle \partial_\lambda \nu_\lambda^R, \eta_\lambda^R \rangle + \langle \nu_\lambda^R, \partial_\lambda \eta_\lambda^R \rangle) + \langle \nu_\lambda^R, (\partial_\lambda \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle \\ &= \mu_\lambda^R \partial_\lambda \langle \nu_\lambda^R, \eta_\lambda^R \rangle + \langle \nu_\lambda^R, (\partial_\lambda \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle \end{aligned}$$

Eq. (4.14) gives $\partial_\lambda \langle \nu_\lambda^R, \eta_\lambda^R \rangle = 0$, and therefore $\partial_\lambda \mu_\lambda^R = \langle \nu_\lambda^R, (\partial_\lambda \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle$, i.e.,

$$\partial_\lambda \mu_\lambda^R = - \int_0^R C_{(0,y)} \mathbb{E}_{(0,y)} [\eta_\lambda^R(Z_R) T_R e^{-\lambda T_R} \mathbb{1}_{Z_R \leq R}] \nu_\lambda^R(dy) \quad (4.17)$$

Since all the integrands are non-negative we have $\partial_\lambda \mu_\lambda^R < 0$. So $\lambda \mapsto \mu_\lambda^R$ is a continuous strictly-decreasing function. Moreover, doing $\lambda = 0$, integrating Eq. (4.13), using Fubini's theorem to integrate first in the z variable, and using Assumption 3.3-(iii), we obtain

$$\begin{aligned} \mu_0^R &= \int_0^R \mathcal{J}_0^R \nu_0^R(dz) \\ &= \int_0^R \int_0^R \int_0^\infty \left(k(\varphi^t(0, y), z) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(0, y)) dt \nu_0^R(dy) dz \\ &= \int_0^R \int_0^\infty \left(\int_0^\infty k(\varphi^t(0, y), z) dz \right) \psi(t|(0, y)) dt \nu_0^R(dy) > 1 \end{aligned}$$

On the other hand, doing $\lambda \rightarrow \infty$, passing to the limit under the expectation of Eq. (4.11) we get for every $f \in C([0, R])$, $\mathcal{G}_\lambda^R f \rightarrow 0$ uniformly as $\lambda \rightarrow \infty$. In particular, by the equicontinuity of \mathcal{G}_λ^R , for every $\delta \in (0, 2)$, there must be λ_* large enough such that for every $f \in C([0, R])$, $\mathcal{G}_\lambda^R f \leq \delta$ for all $\lambda \geq \lambda_*$ and hereby, $\mu_\lambda^R \leq \delta$ for all $\lambda \geq \lambda_*$. Therefore $\mu_\lambda^R \rightarrow 0$ as $\lambda \rightarrow \infty$. In consequence, there must be a unique $\lambda_0 > 0$ such that $\mu_{\lambda_0}^R = 1$. We then define λ_R as the only $\lambda_0 > 0$ such that $\mu_{\lambda_0}^R = 1$ and denote $\eta_R = \eta_{\lambda_R}^R$ the respective eigenfunction. Next, we construct a sequence of h_R from η_R which are to converge to the solution of the intial eigenproblem and we show that we can establish an uniform bound over λ_R .

Step 4 : Construction of h_R .

We extend the definition of K_λ to all $(a, y) \in \mathcal{X}, z \in [0, R]$. Define

$$K_\lambda^R(a, y, z) := \int_0^\infty \left(k(\varphi^t(a, y), z) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(a, y)) e^{-\lambda t} dt,$$

and let

$$h_R(a, y) := \int_0^R \eta_R(z) K_{\lambda_R}^R(a, y, z) dz. \quad (4.18)$$

Hence, taking $a = 0$, since η_R solves Eq. (4.12) for $\mu_\lambda^R = 1$, we have that:

$$h_R(0, y) = \int_0^R \eta_R(z) K_{\lambda_R}^R(0, y, z) dz = \mathcal{G}_{\lambda_R}^R \eta_R(y) = \eta_R(y),$$

and therefore h_R verifies

$$\begin{cases} h_R(\mathbf{x}) = \int_0^R h_R(0, z) K_{\lambda_R}^R(\mathbf{x}, z) dz = C_{\mathbf{x}} \mathbb{E}_{\mathbf{x}} [\eta_R(Z_R) e^{-\lambda_R T_R} \mathbb{1}_{Z_R \leq R}] & \forall \mathbf{x} \in \mathcal{X} \\ h_R(0, y) = \eta_R(y) & \forall y \in (0, R) \end{cases} \quad (4.19)$$

where $C_{\mathbf{x}} = \|K_0(\mathbf{x}, \cdot)\|_{L_1(\mathbb{R}_+)}$. Then, we can repeat the steps of the proof of Lemma 4.2 to show that the truncated renewal equation (4.19) (which is the truncated version of Eq. (4.2)) is equivalent to have the boundary condition

$$\lim_{t \rightarrow +\infty} h_R(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds - \lambda_R t\right) = 0 \quad (4.20)$$

and to have that h_R is solution to the truncated eigenvalue problem

$$\mathcal{Q}_R h_R(a, y) = \lambda_R h_R(a, y)$$

where

$$\begin{aligned} \mathcal{Q}_R h(a, y) &= g(a, y)^\top \nabla h(a, y) \\ &+ \beta(a, y) \left(\int_0^R h(0, z) \left(k(a, y, z) + \frac{\int_R^\infty k(a, y, \zeta) d\zeta}{R} \right) dz - h(a, y) \right). \end{aligned}$$

Hence, developing $\mathcal{Q}_R h_R(0, y)$ one obtains

$$\begin{aligned} \mathcal{Q}_R h_R(0, y) &= g_1(0, y) \partial_a h_R(0, y) + g_2(0, y) \partial_y h_R(0, y) \\ &+ \beta(0, y) \left(\int_0^R h_R(0, z) \left(k(a, y, z) + \frac{\int_R^\infty k(a, y, \zeta) d\zeta}{R} \right) dz - h_R(a, y) \right). \end{aligned}$$

Therefore $\eta_R = h_R(0, \cdot)$ is solution to

$$\begin{aligned} \lambda_R \eta_R(y) &= g_2(0, y) \eta_R'(y) - \beta(0, y) \eta_R(y) \\ &+ \beta(0, y) \int_0^R \eta_R(z) \left(k(0, y, z) + \frac{\int_R^\infty k(a, y, \zeta) d\zeta}{R} + g_1(0, y) \partial_a K_{\lambda_R}^R(0, y, z) \right) dz \end{aligned} \quad (4.21)$$

In our case, Assumption 3.3-(ii) which imposes $\beta(0, y) = 0$ for every initial size y simplifies this last equation into

$$g_2(0, y) \eta_R'(y) - \lambda_R \eta_R(y) = 0$$

Therefore for all $R > 1$, if we impose the normalisation condition $\eta_R(1) = 1$, we have

$$\eta_R(y) = \exp\left(\lambda_R \int_1^y \frac{1}{g_2(0, z)} dz\right), \quad y \in [0, R] \quad (4.22)$$

Finally, coming back to (4.11) and (4.12), we have for all $y \in (0, R)$,

$$\begin{aligned} \eta_R(y) &= C_{(0, y)} \mathbb{E}_{(0, y)} [\eta_R(Z_R) e^{-\lambda T_R} \mathbb{1}_{Z_R \leq R}] \\ \iff 1 &= C_{(0, y)} \mathbb{E}_{(0, y)} \left[\frac{\eta_R(Z_R)}{\eta_R(y)} e^{-\lambda T_R} \mathbb{1}_{Z_R \leq R} \right] \\ \iff 1 &= C_{(0, y)} \mathbb{E}_{(0, y)} \left[\exp\left(\lambda \left(\int_y^{Z_R} \frac{1}{g_2(0, z)} dz - T_R \right)\right) \right] \end{aligned}$$

In particular the last equation characterises λ_R as the unique $\lambda > 0$ such that for all $y \in (0, R)$, the following Euler-Lotka-type equation is verified

$$1 = C_{(0,y)} \mathbb{E}_{(0,y)} \left[\exp \left(\lambda \left(\int_y^{Z_R} \frac{1}{g_2(0,z)} dz - T_R \right) \right) \right]. \quad (4.23)$$

Step 5 : Uniform bound for λ_R (Fragmentation case)

Suppose that for all $a \geq 0$, $\text{supp } k(a, y, \cdot) \subseteq (0, y)$. This is, the newborns sizes are almost surely smaller than the parent size. Hence, for all initial size y we have

$$\mathbb{P}_{(0,y)} \left(T_R > \int_y^{Z_R} \frac{1}{g_2(A_{0,y}(z), z)} dz \right) = 1. \quad (4.24)$$

Indeed, from Lemma 3.1-(4.) we have that $\phi_{0,y}^{-1}(A_{0,y}(z), z) = \int_y^z \frac{1}{g_2(A_{0,y}(\zeta), \zeta)} d\zeta$ is the time needed to go from size y to z following the deterministic flow only, and it has to be smaller than the division time at which the trajectory jumps to z . Then, thanks to Assumption 3.3-(i) which gives $g_2(0, y) \geq g_2(a, y)$, we have also that

$$\mathbb{P}_{(0,y)} \left(T_R > \int_y^{Z_R} \frac{1}{g_2(0, z)} dz \right) = 1.$$

Therefore for all $\lambda > 0$

$$\exp \left(\lambda \left(\int_y^{Z_R} \frac{1}{g_2(0, z)} dz - T_R \right) \right) \leq 1 \in L^1(\mathbb{R}_+^2, p_{(0,y)} dt dz), \quad \mathbb{P}_{(0,y)}\text{-a.s.}, \quad (4.25)$$

and by dominated convergence if λ_R converges to $+\infty$ as $R \rightarrow \infty$, then

$$\mathbb{E}_{(0,y)} \left[\exp \left(\lambda_R \left(\int_y^{Z_R} \frac{1}{g_2(0, z)} dz - T_R \right) \right) \right] \rightarrow 0$$

which contradicts Eq. (4.23). So there must exist $\bar{\Lambda} > 0$ such that for all $R > 1$, $\lambda_R < \bar{\Lambda}$. Moreover, analogous to Step 3, if we differentiate Eq. (4.15) in the sense of distributions with respect to R , we obtain

$$\partial_R \mu_\lambda^R = \langle \nu_\lambda^R, (\partial_R \mathcal{G}_\lambda^R) \eta_\lambda^R \rangle.$$

Again, the definition \mathcal{G}_λ^R gives us that this derivative can be computed in the strong sense. Indeed, for any positive continuous function $f : [0, R] \rightarrow \mathbb{R}_+$ we have

$$\begin{aligned} & \partial_R \mathcal{G}_\lambda^R f(y) \\ &= \frac{\partial}{\partial R} \int_0^R f(z) \int_0^\infty \left(k(\varphi^t(0, y), z) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(0, y)) e^{-\lambda t} dt dz \\ &= f(R) \left(\int_0^\infty \left(k(\varphi^t(0, y), R) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(0, y)) e^{-\lambda t} dt \right) \\ &\quad - \int_0^R f(z) \int_0^\infty \frac{1}{R} \left(k(\varphi^t(0, y), R) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(0, y)) e^{-\lambda t} dt dz \\ &= \frac{1}{R} \int_0^R (f(R) - f(z)) \left(\int_0^\infty \left(k(\varphi^t(0, y), R) + \frac{\int_R^\infty k(\varphi^t(0, y), \zeta) d\zeta}{R} \right) \psi(t|(0, y)) e^{-\lambda t} dt \right) dz \end{aligned}$$

which is positive whenever f is an increasing function. Since Eq. (4.22) gives that for every fixed λ , $\eta_\lambda^R(y)$ is increasing in y , then $(\partial_R \mathcal{G}_\lambda^R) \eta_\lambda^R > 0$ and therefore $\partial_R \mu_\lambda^R > 0$. In particular, the sequence of λ_R , which is defined as the values of λ such that $\mu_\lambda^R = 1$, is then also increasing in R .

Step 6 : Identification of the limit

Step 5 gives that $(\lambda_R)_R$ is an increasing bounded sequence as $R \rightarrow \infty$, so with a limit written $\lambda > 0$. Moreover, for each λ_R exists a unique h_R associated, defined by Eq. (4.18). The family of h_R is equibounded and equicontinuous thanks to Eq. (4.20), Eq. (4.22) and the bound on λ_R . Note indeed that Eq. (4.22) depends on R only through λ_R . We can therefore extract a subsequence converging to some (λ, h) as $R \rightarrow \infty$. We must now check that (λ, h) is a good pair of eigenelements, which we do by dominated convergence. In Step 4 we have constructed h_R such that it is solution to Equations (4.19) and (4.20) which we repeat below to justify each limit.

$$\begin{cases} h_R(\mathbf{x}) = C_{\mathbf{x}} \mathbb{E}_{\mathbf{x}} [\eta_R(Z_R) e^{-\lambda_R T_R} \mathbb{1}_{Z_R \leq R}] & \forall \mathbf{x} \in \mathcal{X} \\ h_R(0, y) = \eta_R(y) & \forall y \in (0, R) \\ h_R(\varphi^t(\mathbf{x})) \underset{t \rightarrow \infty}{\sim} \exp\left(\int_0^t \beta(\varphi^s(\mathbf{x})) ds + \lambda_R t\right) \end{cases}$$

The normalisation constant $C_{\mathbf{x}}$ is already the one required in the limit case. For the expectation term, recalling from Eq. (4.22) that

$$\frac{\eta_R(y_2)}{\eta_R(y_1)} = \exp\left(\lambda_R \int_{y_1}^{y_2} \frac{1}{g_2(0, z)} dz\right)$$

and using Eq. (4.25) in Step 5, we deduce that for all $y \in (0, R)$,

$$\eta_R(Z_R) e^{-\lambda_R T_R} \leq \eta_R(y) \quad \mathbb{P}_{(0, y)\text{-a.s.}}$$

Therefore for all $R > 1$,

$$\mathbb{E}_{(a, y)} [\eta_R(Z_R) e^{-\lambda_R T_R} \mathbb{1}_{Z_R \leq R}] \leq \eta_R(y) < +\infty$$

and we can pass to the limit under the expectations and conclude that the limit h and λ verify the renewal formula

$$h(\mathbf{x}) = C_{\mathbf{x}} \mathbb{E}_{\mathbf{x}} [h(0, Z) e^{-\lambda T}] \quad \forall \mathbf{x} \in \mathcal{X}$$

which is Eq. (4.4) and is equivalent to Eq. (4.2). Thus, by Lemma 4.2 the couple (λ, h) is almost everywhere solution to $\mathcal{Q}h = \lambda h$.

□

Remark 4.7. The assumption $\beta(0, \cdot) \equiv 0$ is crucial for the characterisation of h in Step 4 of the proof of Proposition 4.1. The case $\beta(0, x) > 0$ could possibly be treated, but it would require additional assumptions in order to have $a \mapsto K_\lambda^R(a, x, z) \in W_{loc}^{1,1}(\mathbb{R}_+)$ and to then control the age derivatives of the kernel K_λ^R . Then, Eq. (4.21) would be a scalar transport equation for h_λ^R , which thereby admits an elliptic maximum principle. Nonetheless, the assumption $\beta(0, \cdot) \equiv 0$, while being perfectly biologically meaningful, allows us to avoid this technicalities.

5 Doeblin minoration for compacts

We want to prove the following Doeblin petite-set condition for all the compact sets of \mathcal{X} .

Proposition 5.1. *Let P_t be the Markov process characterised by the infinitesimal generator \mathcal{A} defined by Eq. (2.5). If Assumptions 3.3 are verified, then every compact $\mathcal{K} \subset \mathbb{R}_+^2$ is a petite-set for some skeleton chain of P_t . This is, there is a non-trivial discrete sampling measure μ over \mathbb{R}_+ and a non-trivial measure ν over \mathbb{R}_+^2 such that*

$$\langle \mu, \delta_{\mathbf{x}} P_t f \rangle = \int_0^\infty P_t f(\mathbf{x}) \mu(dt) \geq \langle \nu, f \rangle \quad \forall \mathbf{x} \in \mathcal{K}$$

Before the proof we will introduce some useful lemmas. First, we recall Duhamel formula (5.1), which describes the trajectories driven by the semigroup P_t and allows us to extend the definition of the semigroup as the mild solution of an iterative evolution equation.

Lemma 5.2 (Duhamel formula). *For all $\mathbf{x} \in \mathcal{X}$, $f \in C_b^{1,1}(\mathcal{X})$, P_t is the mild solution to*

$$\begin{aligned} P_t f(\mathbf{x}) &= f(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds\right) \\ &+ \int_0^t \psi(s|\mathbf{x}) \int_0^\infty P_{t-s} f(0, z) \frac{h(0, z) k(\varphi^s(\mathbf{x}), z)}{\int_0^\infty h(0, z') k(\varphi^s(\mathbf{x}), z') dz'} dz ds, \end{aligned} \quad (5.1)$$

Proof. A classical probabilistic proof consists in writing $P_t f(\mathbf{x})$ conditionally to the occurrence of the first jump. It is also possible to prove it by means of a variation of parameters method, as in Corollary 1.7 from [12], for example. Here we provide the probabilistic proof. Let X a Markov process whose law is given by generator \mathcal{A} defined in Eq. (2.5). Recall from definition 3.2 the random variables T and Z which represent the time of the first jump and the new size after the first jump. Note however that the transition kernel of the Markovian generator \mathcal{A} has been rescaled, so that the joint law of (T, Z) under $\mathbb{P}_{\mathbf{x}}$ is from now on given by the density function

$$p_{\mathbf{x}}(t, z) = \psi(t|\mathbf{x}) \cdot \frac{\beta(\mathbf{x}) \frac{h(0, z)}{h(\mathbf{x})} k(\varphi^t(\mathbf{x}), z)}{\int_0^\infty \beta(\mathbf{x}) \frac{h(0, z')}{h(\mathbf{x})} k(\varphi^t(\mathbf{x}), z') dz'} = \psi(t|\mathbf{x}) \cdot \frac{h(0, z) k(\varphi^t(\mathbf{x}), z)}{\int_0^\infty h(0, z') k(\varphi^t(\mathbf{x}), z') dz'}$$

where the probability density of the transition $\mathbf{x} \mapsto (0, z)$ is computed as the ratio between the transition rate of $\mathbf{x} \mapsto (0, z)$ and the total transition rate, as described by the generator \mathcal{A} . Hence, by conditioning on T under $\mathbb{P}_{\mathbf{x}}$ and using the strong Markov property of X , we have:

$$\begin{aligned} P_t f(\mathbf{x}) &= \mathbb{E}_{\mathbf{x}} [f(X_t)] = \mathbb{E}_{\mathbf{x}} [f(X_t) \mathbf{1}_{T>t}] + \mathbb{E}_{\mathbf{x}} [f(X_t) \mathbf{1}_{T \leq t}] \\ &= \mathbb{E}_{\mathbf{x}} [f(X_t) | T > t] \mathbb{P}_{\mathbf{x}}(T > t) + \mathbb{E}_{\mathbf{x}} [\mathbb{E}_{\mathbf{x}} [f(X_t) | T] \mathbf{1}_{T \leq t}] \\ &= \mathbb{E}_{\mathbf{x}} [f(X_t) | T > t] \mathbb{P}_{\mathbf{x}}(T > t) + \mathbb{E}_{\mathbf{x}} [\mathbb{E}_Z [f(X_{t-T})] \mathbf{1}_{T \leq t}] \\ &= f(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds\right) \\ &+ \int_0^t \psi(s|\mathbf{x}) \int_0^\infty P_{t-s} f(0, z) \frac{h(0, z) k(\varphi^s(\mathbf{x}), z)}{\int_0^\infty h(0, z') k(\varphi^s(\mathbf{x}), z') dz'} dz ds. \end{aligned}$$

□

We can give now the proof of Proposition 5.1:

Proof of Proposition 5.1. Let $\mathbf{x} \in \mathcal{X}$ compact such that $\mathcal{X} \subset [\underline{a}, \bar{a}] \times [\underline{y}, \bar{y}]$. We iterate once Duhamel's formula (5.1), using the positivity of P_t :

$$\begin{aligned}
P_t f(\mathbf{x}) &= f(\varphi^t(\mathbf{x})) \exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds\right) \\
&\quad + \int_0^t \psi(s|\mathbf{x}) \int_0^\infty \frac{h(0, z)k(\varphi^s(\mathbf{x}), z)}{\int_0^\infty h(0, z')k(\varphi^s(\mathbf{x}), z') dz'} \left\{ f(\varphi^{t-s}(0, z)) \exp\left(-\int_0^{t-s} \beta(\varphi^s((0, z))) ds\right) \right. \\
&\quad \left. + \int_0^{t-s} \psi(u|\mathbf{x}) \int_0^\infty P_{t-s-u} f(0, \xi) \frac{h(0, \xi)k(\varphi^u(0, z), \xi)}{\int_0^\infty h(0, \xi')k(\varphi^u(0, z), \xi') d\xi'} d\xi du \right\} dz ds \\
&\geq \int_0^t \psi(s|\mathbf{x}) \int_0^\infty f(\varphi^{t-s}(0, z)) \exp\left(-\int_0^{t-s} \beta(\varphi^s((0, z))) ds\right) \frac{h(0, z)k(\varphi^s(\mathbf{x}), z)}{\int_0^\infty h(0, z')k(\varphi^s(\mathbf{x}), z') dz'} dz ds
\end{aligned} \tag{5.2}$$

To obtain the desired result we aim to solve two crucial steps:

- i. First, to prove the existence of some C^1 -diffeomorphism which could allow us to change variables inside the latter integral as to obtain a measure over \mathcal{X} .
- ii. Second, to bound from below the resulting integral uniformly for every $\mathbf{x} \in \mathcal{X}$, using its compactness.

Fix some final time $t \geq 0$, and define $\gamma_t : \mathcal{X} \rightarrow \mathcal{X}$ as

$$\gamma_t(s, z) := \varphi^{t-s}(0, z).$$

We show first that it's a differentiable function. Fix s, z and suppose

$$(a, y) = \gamma_t(s, z).$$

Then the function u defined as $u(s) = \gamma_t(s, z)$ is the unique solution to the Initial Value Problem

$$\begin{cases} u'(s) = -g(u(s)), & s \leq t \\ u(0) = (a, y) \end{cases}$$

Thus, $\partial_s \gamma_t(s, z) = -g(u(s))$. Moreover, by Lemma 3.1, the smoothness of the vector field g and the fact that the ODE system is autonomous gives the smoothness of the flow with respect to the initial condition. Thus, the Jacobian matrix of γ_t equals for all $s \leq t$ and $z > 0$:

$$\mathcal{D}\gamma_t(s, z) = \begin{bmatrix} -g(\varphi^{t-s}(0, z)) & \partial_z \varphi^{t-s}(0, z) \end{bmatrix}, \tag{5.3}$$

where, from Lemma 3.1-2, the derivative of the flow with respect to the initial size is given by

$$\partial_z \varphi^t(0, z) = \exp\left(\int_0^t \mathcal{D}g(\varphi^s(0, z)) ds\right) \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

where we recall that $\mathcal{D}g$ stands for the Jacobian matrix of g and $\exp(\cdot)$ is an exponential matrix. Moreover, let $r \mapsto Y_{(a,y)}(r)$ be the unique orbit of the vector field g passing through the point (a, y) . Its is straightforward that $z = Y_{(a,y)}(0)$, so that the inverse of γ_t is given for all $(a, y) \in \mathbb{R}_+^2$ by

$$\gamma_t^{-1}(a, y) = \left(t - \phi_{0, Y_{(a,y)}(0)}^{-1}(a, y), Y_{(a,y)}(0)\right).$$

Fig. 2 summarises graphically the change of variables and the definition of γ_t^{-1} . Given a, y, \mathbf{x} and t , the inversion of γ consists in determinating the value of ordinate z when




Figure 2: Graphical description of the change of variables defined by γ_t

the integral curve flowing towards (a, y) hits the y -axis and the time $t - s$ required to go from this point to (a, y) . Since $Y_{(a,y)}$ (green line) is known, the inversion is direct. We conclude that γ_t is a C^1 -diffeomorphism and then performing the change of variables $(a, y) = \gamma_t(s, z)$ in the RHS of Eq. (5.2) gives

$$\begin{aligned}
 P_t f(\mathbf{x}) \geq \int_{\mathbb{R}_+^2} f(a, y) & \left\{ \psi \left(t - \phi_{0, Y_{(a,y)}(0)}^{-1}(a, y) | \mathbf{x} \right) \right. \\
 & \exp \left(- \int_0^{\phi_{0, Y_{(a,y)}(0)}^{-1}(a, y)} \beta \left(\varphi^s \left((0, Y_{(a,y)}(0)) \right) \right) ds \right) \\
 & \frac{h(0, Y_{(a,y)}(0)) k \left(\varphi^{t - \phi_{0, Y_{(a,y)}(0)}^{-1}(a, y)}(\mathbf{x}), Y_{(a,y)}(0) \right)}{\int_0^\infty h(0, z) k \left(\varphi^{t - \phi_{0, Y_{(a,y)}(0)}^{-1}(a, y)}(\mathbf{x}), z \right) dz} \\
 & \left. \frac{1}{|\det \mathcal{D}\gamma_t(\gamma_t^{-1}(a, y))|} \mathbf{1}_{\phi_{0, Y_{(a,y)}(0)}^{-1}(a, y) \leq t} \right\} da dy. \quad (5.4)
 \end{aligned}$$

Now, using Assumptions 3.3, we can bound the functions and the Jacobian found in the obtained integral. First, since $g \geq 0$, note that $\|\varphi^t(\mathbf{x})\| \geq \|\varphi^s(\mathbf{x})\|$ for all $t > s$. Second, $\beta_- g_1(\mathbf{x}) \leq \beta(\mathbf{x}) \leq \beta_+ g_1(\mathbf{x})$. And third, by the definition of the flow, $\int_0^t g_1(\varphi^s(\mathbf{x})) ds = \varphi_1^t(\mathbf{x})$ which equals the age at time t of an individual with trait \mathbf{x} at time 0. Then, recalling that $\mathcal{K} \subset [\underline{a}, \bar{a}] \times [\underline{y}, \bar{y}]$, for all $t > 0$ we obtain the following bounds:

- i. For all $(a_0, y_0) \in \mathcal{K}$, using the superior bounds on g_1 from Assumptions 3.3-i. we have

$$\varphi_1^t(a_0, y_0) = a_0 + \int_0^t g_1(\varphi^s(a_0, y_0)) ds \leq a_0 + \int_0^t c_1(1 + \varphi_1^s(a_0, y_0)) ds$$

Hence, by Gronwall inequality

$$\varphi_1^t(a_0, y_0) \leq (a_0 + c_1 t) e^{c_1 t} \leq (\bar{a} + c_1 t) e^{c_1 t}$$

Analogously, using the lower bounds on g_1 from Assumptions 3.3-i., we obtain

$$\varphi_1^t(a_0, y_0) \geq a_0 e^{c_0 t} \geq \underline{a} e^{c_0 t}$$

ii. From the previous result, for all $\mathbf{x} \in \mathcal{X}$

$$\exp\left(-\int_0^t \beta(\varphi^s(\mathbf{x})) ds\right) \geq \exp\left(-\beta_+ \int_0^t g_1(\varphi^s(\mathbf{x})) ds\right) = e^{-\beta_+ \varphi_1^t(\mathbf{x})} \geq e^{-\beta_+ (\bar{a} + c_1 t) e^{c_1 t}},$$

iii. Analogously

$$\beta(\varphi^t(\mathbf{x})) \geq \beta_- g_1(\varphi^t(\mathbf{x})) \geq \beta_- c_0 \varphi_1^t(\mathbf{x}) \geq \beta_- c_0 \underline{a} e^{c_0 t}.$$

Therefore there are some constants $A_0, B_0 > 0$ such that

$$\psi(t-s|\mathbf{x}) \geq A_0 \exp\left(-B_0(1+t-s)e^{c_1(t-s)}\right) \quad (5.5)$$

iv. Moreover, recall that the eigenfunction h is solution to Eq. (4.2). Then, by Fubini's Theorem, for all $\mathbf{x} \in \mathcal{X}$,

$$h(\mathbf{x}) = \int_0^\infty \left(\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \right) \psi(t|\mathbf{x}) e^{-\lambda t} dt.$$

Thus, in particular, the integrability gives us that

$$\left(\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \right) \frac{\psi(t|\mathbf{x}) e^{-\lambda t}}{h(\mathbf{x})} \rightarrow 0 \quad \text{as } t \rightarrow +\infty.$$

Therefore, there exist some constants $C_1, C_2 > 0$ such that for all $\mathbf{x} \in \mathcal{X}$ there is some time $T(\mathbf{x}) > 0$ such that for all $t > 0$ we have

$$\left(\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \right) \frac{\psi(t|\mathbf{x}) e^{-\lambda t}}{h(\mathbf{x})} \leq C_1 \mathbf{1}_{t \geq T(\mathbf{x})} + C_2 \mathbf{1}_{t < T(\mathbf{x})}$$

where

$$\sup_{\mathbf{x} \in \mathcal{X}} \sup_{t < T(\mathbf{x})} \left(\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz \right) \frac{\psi(t|\mathbf{x}) e^{-\lambda t}}{h(\mathbf{x})} \leq C_2,$$

since the suprema are taken in a compact set and for a continuous locally bounded function. Then, taking $C_0 = \max\{C_1, C_2\}$ we have

$$\frac{1}{\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz} \geq \frac{\psi(t|\mathbf{x}) e^{-\lambda t}}{C_0 h(\mathbf{x})}$$

where $\psi(t|\mathbf{x})$ can again be bounded by below using Eq. (5.5). Moreover, the continuity of h implies that h is locally bounded and hence, for all $\mathbf{x} \in \mathcal{X}$, $h(\mathbf{x}) \leq H_0 < \infty$. Hence we obtain finally

$$\frac{1}{\int_0^\infty h(0, z) k(\varphi^t(\mathbf{x}), z) dz} \geq \frac{A_0}{C_0 H_0} \exp\left(-B_0(1+t)e^{c_1 t} - \lambda t\right) \quad (5.6)$$

Note that these three estimates give bounds which are dependent only on t .

v. From (5.3), for all $s \leq t$ and $z > 0$, the Jacobian determinant equals

$$\det \mathcal{D}\gamma_t(s, z) = \|g(\varphi^{t-s}(0, z))\| \|\nabla_z \varphi^{t-s}(0, z)\| \sin \theta(s, t, z) \quad (5.7)$$

where $\theta(s, t, z)$ is the angle between $g(\varphi^{t-s}(0, z))$ and $\nabla_z \varphi^{t-s}(0, z)$. Hence, from Lemma 3.1.2, we get

$$|\det \mathcal{D}\gamma_t(s, z)| \leq \|g(\varphi^{t-s}(0, z))\| \left\| \exp \left(\int_0^{t-s} \mathcal{D}g(\varphi^u(0, z)) du \right) \right\|,$$

where $\|\cdot\|$ is the matrix norm induced by $\|\cdot\|$, and therefore

$$\begin{aligned} |\det \mathcal{D}\gamma_t(\gamma_t^{-1}(a, y))| &\leq \|g(a, y)\| \left\| \exp \left(\int_0^{\phi_{0, Y(a, y)}^{-1}(a, y)} \mathcal{D}g(\varphi^u(0, Y(a, y)(0))) du \right) \right\| \\ &=: \|g(a, y)\| E_0(a, y), \end{aligned} \quad (5.8)$$

Note that this bound depends only on (a, y) and neither on \mathbf{x} or t .

Hence, coming back to Eq. (5.4) and applying the bounds (5.5), (5.6) and (5.6) to the integrands, we obtain

$$\begin{aligned} P_t f(\mathbf{x}) &\geq \int_{\mathbb{R}_+^2} f(a, y) \left\{ \frac{A_0^2}{C_0 H_0} \exp(-2B_0(1+t)e^{c_1 t} - \lambda t) \right. \\ &\quad \exp \left(- \int_0^{\phi_{0, Y(a, y)}^{-1}(a, y)} \beta(\varphi^s(0, Y(a, y)(0))) ds \right) \\ &\quad \left. h(0, Y(a, y)(0)) k \left(\varphi^{t-\phi_{0, Y(a, y)}^{-1}(a, y)}(\mathbf{x}), Y(a, y)(0) \right) \right. \\ &\quad \left. \frac{1}{\|g(a, y)\| E_0(a, y)} \mathbb{1}_{\phi_{0, Y(a, y)}^{-1}(a, y) \leq t} \right\} da dy. \end{aligned} \quad (5.9)$$

Now, we make use of the petite-set condition which allows us to average the value of $P_t f(\mathbf{x})$ against a discrete sampling measure $\mu(dt)$ over a Δ -skeleton. This is, consider some $\Delta > 0$, which will be fixed later on, and a measure μ over $\{j\Delta : j \in \mathbb{N}\}$, characterised by a sequence $(\mu_j)_{j \in \mathbb{N}}$ with $\sum \mu_j = 1$ and $\mu_j > 0$ for all $j \in \mathbb{N}$. We have

$$\begin{aligned} \langle \mu, \delta_{\mathbf{x}} P \cdot f \rangle &\geq \sum_{j=0}^{\infty} \mu_j \int_{\mathcal{X}} f(a, y) k \left(\varphi^{j\Delta - \phi_{0, Y(a, y)}^{-1}(a, y)}(\mathbf{x}), Y(a, y)(0) \right) \\ &\quad \zeta(a, y) e^{-\tilde{\beta} j \Delta} e^{j \Delta} \mathbb{1}_{\phi_{0, Y(a, y)}^{-1}(a, y) \leq j \Delta} da dy, \end{aligned}$$

where the the function $\zeta(a, y)$ is constructed by regrouping all the terms which depend only on (a, y) (and neither on \mathbf{x} or t), and the constant $\tilde{\beta} > 0$ is obtained after selecting only the dominant term inside the exponential. Now, it remains to loose the dependency on \mathbf{x} using that $\mathbf{x} \in \mathcal{X}$ to find a uniform lower bound for the whole compact. By Assumption 3.3-(iv), we have that for all z , exists $D(z) \subset \mathbb{R}_+$ such that $k(\varphi^s(\mathbf{x}), z) > \varepsilon(z) \mathbb{1}_{D(z)}(\varphi^s(\mathbf{x}))$. Then, let

$$\mathcal{T}(\mathbf{x}, z) := \{s > 0 : \varphi^s(\mathbf{x}) \in D(z)\},$$

then

$$k(\varphi^s(\mathbf{x}), z) > \varepsilon(z) \mathbb{1}_{\mathcal{T}(\mathbf{x}, z)}(s).$$

Now, let $\Delta = \inf_{z>0} \text{diam}(D(z)) > \delta_- > 0$. Then, for all $\mathbf{x} \in \mathcal{X}$ and $z > 0$ there exists $n = n(\mathbf{x}, z) \in \mathbb{N}$ such that $n\Delta \in \mathcal{T}(\mathbf{x}, z)$. Then for all $\mathbf{x} \in \mathcal{X}$ and $z > 0$,

$$\sum_{j=0}^{\infty} \mathbb{1}_{j\Delta \in \mathcal{T}(\mathbf{x}, z)} \geq 1.$$

Moreover, since for all $z > 0$, $\text{diam}(D(z)) < \delta_+$, there exists some j big enough such that the trajectory leaves $D(z)$. In particular, the compactness of \mathcal{X} implies that it exists j^* such that for all $\mathbf{x} \in \mathcal{X}$

$$\mathbb{1}_{j\Delta \in \mathcal{T}(\mathbf{x}, z)} = 0 \quad \forall j \geq j^*.$$

Therefore for any sampling measure $(\mu_j)_j$ we have

$$\sum_{j=0}^{\infty} \mu_j \mathbb{1}_{j\Delta \in \mathcal{T}(\mathbf{x}, z)} \geq \min_{j \leq j^*} \mu_j,$$

and finally for all fixed $\tau > 0$,

$$\begin{aligned} \sum_{j=0}^{\infty} \mu_j e^{-\tilde{\beta} j \Delta e^{j\Delta}} k(\varphi^{j\Delta - \tau}(\mathbf{x}), z) \mathbb{1}_{\tau \leq j\Delta} &\geq \sum_{j=0}^{\infty} \mu_j e^{-\tilde{\beta} j \Delta e^{j\Delta}} \epsilon(z) \mathbb{1}_{j\Delta - \tau \in \mathcal{T}(\mathbf{x}, z)} \\ &\geq \epsilon(z) \min_{j \leq j^*} \mu_j \min_{j \leq j^*} e^{-\tilde{\beta} j \Delta e^{j\Delta}} \end{aligned}$$

from what we can conclude that

$$\langle \mu, \delta_{\mathbf{x}} P.f \rangle \geq \int_{\mathcal{X}} f(a, y) \nu(a, y) da dy$$

with

$$\nu(a, y) = \zeta(a, y) \epsilon(Y_{(a, y)}(0)) e^{-\tilde{\beta} j^* e^{\Delta j^* \Delta}} \min_{j \leq j^*} \mu_j$$

□

Finally, the proof of the main theorem 2.3 is a direct application of Harris Theorem 2.1.

6 Application: Steady-state size distribution of the adder model of bacterial proliferation

We model the dynamics of an age-size-structured population of *E. coli* bacteria as a measure-valued process with values in $\mathcal{M}_p(\mathcal{X})$, the point measures over the state space $\mathcal{X} = \{(a, y) \in \mathbb{R}_+^2 : 0 < a < y, y > 0\}$, where a represents the added size and y the current size of each cell. This is, the age of a cell is given by the difference between its current size and its initial size, i.e., using our notations

$$A_{(a_0, y_0)}(y) = a_0 + (y - y_0).$$

The importance to consider the added size as a structural variable to accurately model the growing dynamics of *E. coli* has been strongly suggested in the recent years by experimental works and statistical analysis [24]. The dynamics are driven by the generator

$$\begin{aligned} \mathcal{Q}f(a, y) &= \lambda y (\partial_a + \partial_y) f(a, y) \\ &\quad + \lambda y B(a) \left(2 \int_0^1 f(0, \rho y) F(\rho) d\rho - f(a, y) \right) - d_0 f(a, y). \end{aligned} \quad (6.1)$$

In our previous notation this translates as $g(a, y) = (\lambda y, \lambda y)$, $\beta(a, y) = \lambda y B(a)$, and $k(a, y, z) = \frac{1}{y} F\left(\frac{z}{y}\right) \mathbb{1}_{z \leq y}$, where F has support in $[0, 1]$. The growth dynamics correspond to an exponential elongation at constant rate $\lambda > 0$. The second term in \mathcal{Q} represents the divisions, which occur at rate $\beta(a, y) = \lambda y B(a)$ where B is a hazard function such that for every individual,

$$\mathbb{P}(\text{Added size at division} \geq a) = \exp\left(-\int_0^a B(s) ds\right).$$

Hence, the jump term reads as follows: a cell of size y and added size a divides at rate $\lambda y B(a)$, and it's replaced by two cells of added size 0 and sizes ρy and $(1 - \rho)y$ respectively, where ρ is randomly distributed following the density F . The third term represents deaths at a constant rate $d_0 > 0$. We assume that:

Assumptions 6.1. Suppose

(A1) There exist $0 < \underline{b} \leq \bar{b} < \infty$ such that for all $a \geq 0$, $\underline{b} < B(a) < \bar{b}$.

(A2) F is a continuous positive function in $[0, 1]$, with connected support. We call for all $k \geq 0$,

$$m_k = \int_0^1 \rho^k F(\rho) d\rho$$

and suppose that $m_0 = 1$, $m_1 = 1/2$ and $m_2 < +\infty$. Note that, since $\rho \in (0, 1)$ almost surely, then for all $k > 0$ we have $m_k \leq m_1 = 1/2$.

(A3) $\lambda > d_0$.

Theorem 6.2. Under Assumptions (A1)-(A3), there is a unique probability measure π^* such that there exist constants $C, \omega > 0$ which verify Eq. (2.6) with $\Lambda = \lambda - d_0$, $h(a, y) = y$ and $V(a, y) = y^{-1} + y$. Moreover π^* admits a density given explicitly by

$$\pi^*(a, y) = \frac{\exp\left(-\int_0^a B(\alpha) d\alpha\right)}{y^2} \eta^*(y - a),$$

where η^* is the unique solution to the fixed point problem

$$\eta^*(y) = 2 \int_0^1 \left\{ \int_0^{\frac{y}{\rho}} \psi\left(\frac{y}{\rho} - z\right) \eta^*(z) dz \right\} F(\rho) d\rho,$$

where $\psi(a) = B(a) \exp\left(-\int_0^a B(\alpha) d\alpha\right)$.

Proof. 1. **Minoration condition.** It is a direct application of Proposition 5.1, since the same hypothesis in Assumptions 3.3 are verified by Assumptions 6.1. Assumption 6.1-(v) requires some attention. Indeed, since F is bounded and with connected support, $k(a, y, z) = \frac{1}{y} F\left(\frac{z}{y}\right) \mathbb{1}_{z \leq y}$ can be lower bounded in the form $k(a, y, z) > \varepsilon(z) \mathbb{1}_{y \in D(z)}$, with $\varepsilon(z)$ of order $1/z$, as represents the example of Fig. 3.




Figure 3: Example of minoration for $k(a, y, z) = \frac{1}{y} F\left(\frac{z}{y}\right) \mathbb{1}_{z \leq y}$ and F given by the probability density function of a Beta distribution. Then we have $k(a, y, z) > \varepsilon(z) \mathbb{1}_{y \in D(z)}$ as required by Assumption 3.3-(v), with $|D(z)| = \delta$ for all z .

In general, we have for all $\delta > 0$:

$$k(a, y, z) > \min_{z' \in [2z, 2z + \delta]} \frac{F(z/z')}{z'} \mathbf{1}_{y \in [2z, 2z + \delta]}$$

and we verify then Assumption 3.3-(v) with $\varepsilon(z) = \min_{z' \in [2z, 2z + \delta]} \frac{F(z/z')}{z'}$ and $D(z) = [2z, 2z + \delta]$ for a chosen $\delta > 0$.

Fig. 4 shows the characteristics curves $y - a = \text{constant}$, and the shadowed region corresponds the space that is a priori reachable from the initial point along trajectories with exactly one jump before time t . It is the version of Fig. 2 in this specific case. Moreover, given an initial point (A in Fig. 2) and total trajectory time, this reachable region is compact, which also simplifies some minorations.

Finally, depending on the choice of the compact set $\mathcal{X} \subset [a, \bar{a}] \times [y, \bar{y}]$ and of δ (which gives also the discretisation timestep of the δ -skeleton), the value of the minorant measure ν can be computed explicitly by numerical approximations, as given in Fig. 5 for different forms of F .




Figure 4: Ideal trajectory from initial point $A = (a_0, y_0)$ to point $D = (a^*, y^*)$ in time t . The individual spends a time $t - s$ growing from A to B . Then, it divides and renews at point C . Finally, it grows the remaining time s until point D .

2. **Lyapunov-Foster condition.** Let $V(a, y) = y^{-1} + y$ with. It is clear that $V(a, y) \rightarrow \infty$ as $|(a, y)| \rightarrow \infty$. Let $v(a, y) = y^k$, then

$$\begin{aligned} \mathcal{A}v(a, y) &= \lambda y^k + 2\lambda y B(a) \int_0^1 (\rho^k y^k - y^k) \rho F(\rho) d\rho \\ &= \left(k\lambda + 2 \left(m_{k+1} - \frac{1}{2} \right) \lambda y B(a) \right) v(a, y) \end{aligned}$$

So, for $V(a, y) = y^{-1} + y$, we obtain

$$\mathcal{A}V(a, y) = -\lambda V(a, y) + \Delta(a, y),$$

where

$$\Delta(a, y) := 2\lambda y + 2\lambda B(a) \left(\left(\frac{1}{2} + \left(m_2 - \frac{1}{2} \right) y^2 \right) \right).$$

We already have $-\lambda < 0$ in the first term of the RHS. It remains to prove that $\Delta(a, y)$ defined in the RHS above, is bounded. Indeed, notice that

$$\Delta(a, y) \leq 2\lambda \left(\bar{b} \left(m_2 - \frac{1}{2} \right) y^2 + y + \frac{\bar{b}}{2} \right)$$

which is quadratic in y with a negative quadratic coefficient since $m_2 - 1/2 \leq 0$. Thus

$$\Delta(a, y) \leq \lambda \left(\bar{b} + \frac{1}{\bar{b}(1 - 2m_2)} \right) =: d \in \mathbb{R}_+ \quad (6.2)$$

So finally we obtain that for every $(a, y) \in \mathbb{R}_+^2$

$$\mathcal{A}V(a, y) \leq -\lambda V(a, y) + d$$

3. Application of Harris' Theorem Using Theorem 2.1 we conclude the existence of some $C, \omega > 0$ such that for every $x \in \mathbb{R}_+^2$ and $t \geq 0$

$$\|\delta_x P_t - \pi\|_{1+V} \leq C(1 + V(x)) \exp(-\omega t) \quad (6.3)$$

Now, using that by construction, $M_t f = e^{\Lambda t} h P_t (f/h)$, we obtain that for all $x \in \mathbb{R}_+^2$,

$$\|e^{-\Lambda t} \delta_x M_t - h(x) \pi^*\|_{1+V} \leq C(1 + V(x)) e^{-\omega t} \quad (6.4)$$

where for every $A \in \mathcal{B}(\mathbb{R}_+^2 \setminus \{0\})$,

$$\pi^*(A) = \int_A \frac{\pi(d\mathbf{x})}{h(\mathbf{x})}$$

Moreover, we know that $h(a, y) = y$. On the other hand, π is the unique solution to $\pi P_t = \pi$, or equivalently, to the dual eigenvalue problem associated to the conservative problem $\pi \mathcal{A} = 0$. From (2.5) we obtain from the latter that π is then the measure solution to the following PDE in the sense of distributions

$$\left\{ \begin{array}{l} (\partial_a + \partial_y)(\lambda y \pi(a, y)) - \lambda y B(a) \pi(a, y) = 0 \\ \pi(0, y) = 2 \int_0^1 \int_0^\infty B(a) \frac{F(\rho)}{\rho} \pi \left(a, \frac{y}{\rho} \right) dad\rho \\ \int_0^\infty \int_0^y \pi(a, y) dady = 1 \end{array} \right. \quad (6.5)$$

We solve it by the method of characteristics. From the first equation of (6.5), π solves the ODE

$$\left\{ \begin{array}{l} \frac{d}{da} \pi(a, y(a)) = - \left(B(a) + \frac{1}{y(a)} \right) \pi(a, y(a)) \\ \pi(0, y(0)) = 2 \int_0^1 \int_0^\infty B(a) \frac{F(\rho)}{\rho} \pi \left(a, \frac{y(0)-a}{\rho} \right) dad\rho \end{array} \right.$$

where the associated characteristics are of the form $y(a) = a + (y(0) - a(0))$. Then, the solution π of (6.5) is given by

$$\pi(a, y) = \frac{\exp \left(- \int_0^a B(\alpha) d\alpha \right)}{y} \eta^*(y - a),$$

where the definition of η^* is inherited from the initial condition of the ODE:

$$\pi(0, y(0)) = \frac{\eta^*(y - a)}{y - a} = 2 \int_0^1 \int_0^\infty B(a) \frac{F(\rho)}{\rho} \pi \left(a, \frac{y(0) - a}{\rho} \right) dad\rho.$$

Note that the RHS still depends implicitly on π . Hence, η^* is solution to the fixed point problem

$$\begin{aligned}\eta^*(x) &= 2 \int_0^1 \int_0^\infty B(a)F(\rho) \exp\left(-\int_0^a B(\alpha)d\alpha\right) \eta^*\left(\frac{x}{\rho} - a\right) dad\rho \\ &= 2 \int_0^1 \int_0^{\frac{x}{\rho}} F(\rho)\psi\left(\frac{x}{\rho} - a\right) \eta^*(a) dad\rho,\end{aligned}$$

where

$$\psi(a) = B(a) \exp\left(-\int_0^a B(\alpha)d\alpha\right)$$

is the probability density function of the added size at division. The existence of a formal solution to this problem is then a by-product of the existence of π , here provided by Harris' Theorem.

Thus finally, the stationary profile of M_t is given by

$$\pi^*(a, y) = \frac{\exp\left(-\int_0^a B(\alpha)d\alpha\right)}{y^2} \eta^*(y - a)$$

□




Figure 5: Minorant measure ν for F given by *i.* the uniform distribution, *ii.* a Beta(5,5) distribution and *iii.* a Beta(20,20) distribution. The values were obtained from numerical approximation using $\underline{y} = 1, \bar{y} = 2$. Only F changes between the three cases.

Remark 6.3. The stability of this model has already been studied in the early works of [16] for an application to plant physiology, and more recently by [15] from which the exponential ergodicity is known. In our case however, the direct application of Harris' Theorem, since the eigenelements of \mathcal{Q} are known, gives a simpler more general argument to obtain the long-time behaviour. More generally, when the drift term $g(a, y)$ is not necessarily given by the exponential elongation, the previous section allows to prove the existence of the suitable eigenelements. This was left as an open question by the works of [15].

Remark 6.4. The proof presented above does not work for singular divisions as given, in lieu of a density F , by ρ distributed according to $\delta_{1/2}(d\rho)$ as in a perfectly symmetric mitosis. Indeed, the change of variables is no longer possible since z would be constant. Moreover, if we try to pursue the method and average in time, one can check that the obtained ν would be the trivial measure for some large enough compacts. Such

a limitation is not really surprising, since the authors of [11] have already shown that if the elongation rate λ is constant for the whole population (as in our case), and the divisions are perfectly symmetrical, then we do not have convergence, and a periodic behaviour is observed.

Remark 6.5. Figure 5 shows the shape of ν for different forms of the density F . As F concentrates we can observe the increasing degeneracy of ν .

Remark 6.6. Here, the existence of η^* is a by-product of the existence and uniqueness of π provided by Harris' Theorem. In contrast, in the works of [15], the existence of the stationary measure depended on the existence of a unique solution to the fixed point problem. Thus, the authors had to show compactness properties of the operator associated to the fixed point problem, which our approach evades. Moreover, our approach allows more general forms for F , which [15] requires to be of compact support strictly included in $]0, 1[$.

Acknowledgments. I thank Sylvie Méléard for her continual guidance and her many readings and suggestions for this manuscript, and Meriem El Karoui which introduced me to the biological motivations behind this model and has also guided me throughout the work. I would also like to thank Bertrand Cloez and Marie Doumic for the rewarding discussions during this work. This work has been supported by the Chair "Modélisation Mathématique et Biodiversité" of Veolia Environnement - École polytechnique - Museum National d'Histoire Naturelle - Fondation X.

References

- [1] Balagué, D., Cañizo, J.A., Gabriel, P.: Fine asymptotics of profiles and relaxation to equilibrium for growth-fragmentation equations with variable drift rates. *Kinetic and Related Models*. 2013, Volume 6, Pages 219-243 **6**, 219 (2013). <https://doi.org/10.3934/KRM.2013.6.219>
- [2] Bansaye, V., Cloez, B., Gabriel, P.: Ergodic behavior of non-conservative semigroups via generalized doebelin's conditions. *Acta Applicandae Mathematicae* **166**(1), 29–72 (2020). <https://doi.org/10.1007/s10440-019-00253-5>
- [3] Bansaye, V., Cloez, B., Gabriel, P., Marguet, A.: A non-conservative harris ergodic theorem. *Journal of the London Mathematical Society* **106**(3), 2459–2510 (2022) <https://londmathsoc.onlinelibrary.wiley.com/doi/pdf/10.1112/jlms.12639>. <https://doi.org/10.1112/jlms.12639>
- [4] Bouguet, S., Florian, J.: A Probabilistic Look at Growth-Fragmentation Equations. *Séminaire de Probabilités XLIX. Lecture Notes in Mathematics* **2215**, 57–74 (2018). https://doi.org/10.1007/978-3-319-92420-5_2
- [5] Canizo, J.A., Gabriel, P., Yoldaş, H.: Spectral gap for the growth-fragmentation equation via harris's theorem. *SIAM Journal on Mathematical Analysis* **53**(5), 5185–5214 (2021) <https://doi.org/10.1137/20M1338654>. <https://doi.org/10.1137/20M1338654>
- [6] Cloez, B., Gabriel, P.: On an irreducibility type condition for the ergodicity of nonconservative semigroups. *Comptes Rendus. Mathématique* **358**(6), 733–742 (2020). <https://doi.org/10.5802/crmath.92>
- [7] Cloez, B., de Saporta, B., Roget, T.: Long-time behavior and darwinian optimality for an asymmetric size-structured branching process. *Journal of Mathematical Biology* **83**(6), 69 (2021). <https://doi.org/10.1007/s00285-021-01695-y>
- [8] Costa, M.: A piecewise deterministic model for a prey-predator community. *The Annals of Applied Probability* **26**(6), 3491–3530 (2016). <https://doi.org/10.1214/16-AAP1182>
- [9] Doumic, M.: Analysis of a population model structured by the cells molecular content. *Math. Model. Nat. Phenom* **2**, 121–152 (2007)
- [10] Doumic Jauffret, M., Gabriel, P.: Eigenelements of a general aggregation-fragmentation model. *Mathematical Models and Methods in Applied Sciences* **20**(05), 757–783 (2010) <https://doi.org/10.1142/S021820251000443X>. <https://doi.org/10.1142/S021820251000443X>

- [11] Doumic, Marie, van Brunt, Bruce: Explicit solution and fine asymptotics for a critical growth-fragmentation equation. *ESAIM: ProcS* **62**, 30–42 (2018). <https://doi.org/10.1051/proc/201862030>
- [12] Engel, K.-J., Nagel, R.: One-parameter semigroups for linear evolution equations. *Semigroup Forum* **63**, 278–280 (1999)
- [13] Fonte, C., Schmutz, V.: Long time behavior of an age- and leaky memory-structured neuronal population equation. *SIAM Journal on Mathematical Analysis* **54**(4), 4721–4756 (2022). <https://doi.org/10.1137/21M1428571>. <https://doi.org/10.1137/21M1428571>
- [14] Gabriel, Pierre: Measure solutions to the conservative renewal equation. *ESAIM: ProcS* **62**, 68–78 (2018). <https://doi.org/10.1051/proc/201862186206>
- [15] Gabriel, P., Martin, H.: Steady distribution of the incremental model for bacteria proliferation. *Networks and Heterogeneous Media* **14**(1), 149–171 (2019). <https://doi.org/10.3934/nhm.2019008>
- [16] Hall, A.J., Wake, G.C., Gandar, P.W.: Steady size distributions for cells in one-dimensional plant tissues. *Journal of Mathematical Biology* **30**, 101–123 (1991)
- [17] Heijmans, H.J.A.M.: The dynamical behaviour of the age-size-distribution of a cell population. In: Metz, J.A.J., Diekmann, O. (eds.) *The Dynamics of Physiologically Structured Populations*, pp. 185–202. Springer, Berlin, Heidelberg (1986)
- [18] Lee, J.M.: *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics. Springer, ??? (2003). <https://doi.org/10.1007/978-1-4419-9982-5>
- [19] Lin, J., Amir, A.: From single-cell variability to population growth. *Physical Review E* **101**(1), 012401 (2020) arXiv:1806.02818. <https://doi.org/10.1103/PhysRevE.101.012401>
- [20] Meyn, S.P., Tweedie, R.L.: Stability of markovian processes iii: Foster-lyapunov criteria for continuous-time processes. *Advances in Applied Probability* **25**, 518–548 (1993)
- [21] Mischler, S., Scher, J.: Spectral analysis of semigroups and growth-fragmentation equations. *Annales de l’Institut Henri Poincaré C, Analyse non linéaire* **33**(3), 849–898 (2016). <https://doi.org/10.1016/J.ANIHPC.2015.01.007>
- [22] Perthame, B.: *Transport Equations in Biology*. Frontiers in Mathematics. Birkhäuser Basel (2007). <https://doi.org/10.1007/978-3-7643-7842-4>.
- [23] Roget, T.: On the long-time behaviour of age and trait structured population dynamics. *Discrete and Continuous Dynamical Systems - B* **24**(6), 2551–2576 (2019). <https://doi.org/10.3934/dcdsb.2018265>
- [24] Taheri-Araghi, S., Bradde, S., Sauls, J.T., Hill, N.S., Levin, P.A., Paulsson, J., Vergassola, M., Jun, S.: Cell-size control and homeostasis in bacteria. *Current Biology* **25**(3), 385–391 (2015). <https://doi.org/10.1016/j.cub.2014.12.009>
- [25] Tomasević, Milica, Bansaye, Vincent, Véber, Amandine: Ergodic behaviour of a multi-type growth-fragmentation process modelling the mycelial network of a filamentous fungus. *ESAIM: PS* **26**, 397–435 (2022). <https://doi.org/10.1051/ps/2022013>
- [26] Torres, N., Perthame, B., Salort, D.: A multiple time renewal equation for neural assemblies with elapsed time model. *Nonlinearity* **35**(10), 5051 (2022). <https://doi.org/10.1088/1361-6544/ac8714>